

**Report on the Peer Consultation Workshop to Discuss a
Proposed Protocol to Assess Asbestos-Related Risk**

Prepared for:

U.S. Environmental Protection Agency
Office of Solid Waste and Emergency Response
Washington, DC 20460

EPA Contract No. 68-C-98-148
Work Assignment 2003-05

Prepared by:

Eastern Research Group, Inc.
110 Hartwell Avenue
Lexington, MA 02421

FINAL REPORT
May 30, 2003

NOTE

This report was prepared by Eastern Research Group, Inc. (ERG), an EPA contractor, as a general record of discussion for the peer consultation workshop on a proposed protocol to assess asbestos-related risk. This report captures the main points of scheduled presentations, highlights discussions among the panelists, and documents the public comments provided at the meeting. This report does not contain a verbatim transcript of all issues discussed, and it does not embellish, interpret, or enlarge upon matters that were incomplete or unclear. EPA will use the information presented during the peer consultation workshop to determine whether the proposed risk assessment methodology can be used to support decisions at asbestos-contaminated sites. Except as specifically noted, no statements in this report represent analyses by or positions of EPA or ERG.

CONTENTS

List of Abbreviations	iii
Executive Summary	v
1. Introduction	1-1
1.1 Background	1-1
1.2 Scope of the Peer Consultation Workshop	1-2
1.2.1 Activities Prior to the Peer Consultation Workshop	1-2
1.2.2 Activities at the Peer Consultation Workshop	1-3
1.2.3 Activities Following the Peer Consultation Workshop	1-4
1.3 Report Organization	1-5
2. Background on the Proposed Protocol to Assess Asbestos-Related Risk	2-1
3. Comments on Topic Area 1: Interpretations of the Epidemiology and Toxicology Literature	3-1
3.1 Lung Cancer	3-1
3.1.1 Lung Cancer and Fiber Type: Inferences from the Epidemiology Literature	3-1
3.1.2 Lung Cancer and Fiber Type: Inferences from Animal Toxicology and Mechanistic Studies	3-6
3.1.3 Lung Cancer and Fiber Dimension: Inferences from the Epidemiology Literature	3-7
3.1.4 Lung Cancer and Fiber Dimension: Inferences from Animal Toxicology and Mechanistic Studies	3-9
3.1.5 Other Issues Related to Lung Cancer	3-10
3.2 Mesothelioma	3-12
3.2.1 Mesothelioma and Fiber Type: Inferences from the Epidemiology Literature	3-12
3.2.2 Mesothelioma and Fiber Type: Inferences from Animal Toxicology and Mechanistic Studies	3-14
3.2.3 Mesothelioma and Fiber Dimension: Inferences from the Epidemiology Literature	3-16
3.2.4 Mesothelioma and Fiber Dimension: Inferences from Animal Toxicology and Mechanistic Studies	3-17
3.3 Exposure Estimates in the Epidemiology Literature	3-18

CONTENTS (Continued)

4.	Comments on Topic Area 2: The Proposed Exposure Index	4-1
4.1	Responses to Charge Question 4	4-1
4.2	Responses to Charge Question 5	4-3
4.3	Responses to Charge Question 6	4-4
5.	Comments on Topic Area 3: General Questions	5-1
5.1	Responses to Charge Question 7	5-1
5.2	Responses to Charge Question 8	5-2
5.3	Responses to Charge Question 9	5-3
5.4	Responses to Charge Question 10	5-4
5.5	Responses to Charge Question 12	5-6
6.	Comments on Topic Area 4: Conclusions and Recommendations	6-1
6.1	Responses to Charge Question 11	6-1
6.2	Development of Final Conclusions and Recommendations	6-6
7.	References	7-1

Appendices

Appendix A	List of Expert Panelists
Appendix B	Premeeting Comments, Alphabetized by Author (includes bios of panelists and the charge to the reviewers)
Appendix C	List of Registered Observers of the Peer Consultation Workshop
Appendix D	Agenda for the Peer Consultation Workshop
Appendix E	Observer Comments Provided at the Peer Consultation Workshop
Appendix F	Observer Post-Meeting Comments

LIST OF ABBREVIATIONS

ATSDR	Agency for Toxic Substances and Disease Registry
EPA	U.S. Environmental Protection Agency
ERG	Eastern Research Group, Inc.
IARC	International Agency for Research on Cancer
IRIS	Integrated Risk Information System
NIOSH	National Institute for Occupational Safety and Health
PCM	phase contrast microscopy
SEM	scanning electron microscopy
SVF	synthetic vitreous fibers
TEM	transmission electron microscopy
µm	micrometers

EXECUTIVE SUMMARY

Eleven expert panelists participated in a peer consultation workshop to review a proposed protocol to assess asbestos-related risks. The protocol is documented in the report, “Technical Support Document for a Protocol to Assess Asbestos-Related Risk, Parts I and II” (Berman and Crump 1999, 2001). At the end of the 2½-day workshop, which was open to the public, the expert panelists drafted the following summary of their findings:

The peer consultation panel strongly endorsed the conceptual approach of developing an updated cancer risk assessment methodology that takes into account fiber type and fiber dimension. The opportunity is at hand to use substantial new information from epidemiology, experimental toxicology, and exposure characterization on what continues to be an extremely important societal issue—assessing the health risks associated with environmental and occupational exposures to asbestos. The panel recommended that EPA proceed in an expeditious manner to consider the panelists’ conclusions and recommendations with a goal of having an updated asbestos risk assessment methodology. It is important that EPA devote sufficient resources so that this important task can be accomplished in a timely and scientifically sound manner. The panel urges that additional analyses underpinning the document, preparation of documentation, and further review be carried out in an open and transparent manner.

Prior to the workshop, the participants received draft copies of the “Methodology for Conducting Risk Assessments at Asbestos Superfund Sites Part 1: Protocol” and “Part 2: Technical Background Document.” The panelists generally found that these documents did not provide a complete and transparent description of how the data were analyzed to support the conclusions presented. The incomplete documentation of methodology precluded the replication of the findings, in advance of the meeting, by several panelists. The methodology used was clarified by the comprehensive presentations that Drs. Berman and Crump made at the workshop. However, future drafts of these documents must

clearly describe the methodologies and include sufficient data, perhaps in appendices, such that the findings can be replicated.

The panelists made the following conclusions and recommendations:

- # **Measurement methods.** Continuing advances have been made in the application of exposure measurement technology for asbestos fibers during the past two decades. These advances include the use of transmission electron microscopy (TEM) and allied techniques (e.g., energy dispersive x-ray detection, or EDS) as an alternative to phase contrast microscopy (PCM), thereby allowing the bivariate (i.e., length and width) characterization of fibers and fiber type. The proposed risk assessment methodology incorporates these advances in the development of an exposure index. The panel was in agreement that this aspect of the new risk assessment methodology represents a substantial advance over the existing methodology.
- # **Integration of exposure and risk assessment models.** A key aspect of the proposed risk assessment methodology is a linking of specific exposure characterization methodology with exposure-response coefficients. It has been emphasized that any change in the exposure characterization metrics must be accompanied by changes in the exposure-response coefficients of the risk assessment models. This was emphasized in the report and the panelists endorsed this view.
- # **Access to additional raw data sets.** The panelists strongly recommended that EPA make every attempt to acquire and analyze raw data sets from key human epidemiological studies. Where possible, it would also be desirable to obtain bivariate (i.e., length and diameter) fiber exposure information for these re-analyses. Several panelists believed that review of additional data sets offers substantial opportunity for improving the proposed risk assessment methodology. In the event that raw data cannot be obtained due to confidentiality reasons or other restrictions, the panelists suggested that the authors consider asking those who have access to the data to conduct the necessary statistical analyses and communicate their results directly to EPA for further consideration.
- # **Fiber diameter.** The proposed risk assessment methodology uses a diameter cut-off of 0.5 micrometers (μm) for considering fibers. The report states that fibers 0.7 μm in diameter can reach the respiratory zone of the lung. A few panel members indicated that the fiber diameter cut-off could be as high as 1.5 μm during oral breathing. The 0.4 μm cut-off came from rat data, but larger diameters would be expected to be respirable in humans. There was general agreement that the diameter cut-off should be between 0.5 and 1.5 μm . This issue is deserving of further analysis.

Fiber length. The Berman and Crump analyses made a significant contribution by obtaining and analyzing membrane filters from the animal inhalation studies in Edinburgh and conducting quality-assured bivariate length and distribution analyses by TEM—thereby greatly reducing the uncertainty of the exposure side of the exposure-response relationship for chronic fiber exposure in rats. Unfortunately, correspondingly detailed information on bivariate size distribution is not available for humans. This leads to the need to use the animal data, although one must always recognize the uncertainties associated with interspecies extrapolations such as anatomic characteristics and respirability between species. Future analyses may benefit from using other available laboratory animal data sets and human data sets.

The fiber length distributions for the human cohort exposures are much more uncertain. For the Wittenoom, Quebec, and South Carolina cohorts, there are limited fiber length distribution data based on TEM analysis from historic membrane filter samples, but only fiber categories longer than 5 μm and longer than 10 μm were counted. For all other cohorts, the measurements were limited to PCM fiber counts for all fibers greater than 5 μm in length in some, and particle counts (10x objective) on midget impinger samples in others. Both methods do not measure thin fibers, do not discriminate between asbestos and other mineral particles, and provide no information on the concentrations of fibers longer than 10, 20, or 40 μm , or inter-laboratory variations in optical resolution and counting rules. As one approach to addressing the varying uncertainty in assessing exposure in the different studies, Berman and Crump used the available information to make adjustments to the uncertainty ranges in the exposure-response coefficients. The workshop panel welcomed this initiative but suggested alternative approaches (see “Methods,” below).

Some panelists felt that an Exposure Assessment Workshop, with participants having a broad range of expertise, could evaluate the uncertainties in historic occupational data sets’ exposure measurements. They felt such a workshop could result in a more confident assessment of exposure-response relationships for populations exposed to a variety of amphiboles, chrysotile, and mixtures. With incorporation of other available knowledge on fiber type, process, smoking (if available), and the relative number of excess lung cancer and mesothelioma, it may well be possible to gain a much clearer understanding of the roles of these variables as causal factors for these asbestos-associated cancers. In addition, the workshop would prove valuable in further discussion of mineralogical, geological, and industrial hygiene issues with regard to application of the model to risk assessment in environmental sites of concern.

The Berman and Crump index assigns zero risk to fibers less than 5 μm in length. Fibers between 5 and 10 μm are assigned a risk that is one three-hundredth of the risk assigned to fibers longer than 10 μm . Panelists agreed that there is a considerably greater risk for lung cancer for fibers longer than 10 μm . However, the panel was uncertain as to an exact cut size for length and the magnitude of the relative potency. The panelists also agreed that the available

data suggest that the risk for fibers less than 5 μm in length is very low and could be zero. This specific issue was addressed by an expert panel convened by the Agency for Toxic Substances and Disease Registry (ATSDR) in October 2002. Some panelists suggested that, for mesothelioma, greater weight should perhaps be assigned to fibers in the 5 to 10 μm length range and to thinner fibers.

- # **Fiber type.** For *mesothelioma*, the panelists supported the use of different relative carcinogenic potencies for different fiber types. The panelists unanimously agreed that the available epidemiology studies provide compelling evidence that the carcinogenic potency of amphibole fibers is two orders of magnitude greater than that for chrysotile fibers. There was some discussion about the precise ratio expressed due to questions about the availability of exposure data in existing studies (e.g., Wittenoom). There was recognition that time since first exposure is an important factor in determining risk for mesothelioma and some discussion is needed on the importance of duration and intensity of exposure.

For *lung cancer*, the panelists had differing opinions on the inferences that can be made on the relative potency of chrysotile and amphibole fibers. Some panelists supported the finding that amphibole fibers are 5 times or more potent for lung cancer than are chrysotile fibers. Other panelists did not think the statistical analyses in the draft methodology document supports this relative potency and wondered if additional review of the epidemiological data might identify factors other than fiber type (e.g., industry considered) that provide further insights on the matter. These other factors can then be considered when the risk assessment is applied.

- # **Cleavage fragments.** The panel knew of little data to directly address the question as to whether cleavage fragments of equal durability and dimension as fibers would have similar or dissimilar potency for lung cancer. The general view is that data indicate that durability and dimension are critical to pulmonary pathogenesis. Therefore, it is prudent at this time to assume equivalent potency for cancer in the absence of other information to the contrary. Consideration of conducting a rat inhalation study using tremolite cleavage fragments was recommended to address this issue. For mesothelioma, it was viewed that thin fibers greater than 5 μm in length are more important. Cleavage fragments that do not meet these criteria would not contribute to risk of mesothelioma.

- # **Other amphiboles.** The panel agreed with the report's conclusion that the potency of currently regulated and unregulated amphibole fibers should be considered equal based on the reasoning that similar durability and dimension would be expected to result in similar pathogenicity.

- # **Methods.** The panelists extensively discussed the approach to conducting the meta-analysis of the large number of epidemiological studies. A number of the panelists urged that consideration be given to using more traditional approaches that would include development and application of specific criteria for inclusion of studies into the exposure-response analysis, examination of

heterogeneity and sources of the heterogeneity, and the use of sensitivity analysis to identify influential studies.

The panelists also urged, in the study-specific analysis, exploration of alternative exposure-response models other than the lung cancer and mesothelioma risk models EPA has been using since 1986. This would possibly include non-linear response models (e.g., log-linear models), examination of separate effects for concentration and duration, time since first exposure, time since cessation of exposure, possibly dropping the “ α factor,” and different methods for measurement error. The adequacy of different models should be examined using goodness of fit statistics across all studies. The possibility of internal analyses should be re-examined (i.e., it may be possible to obtain partial data, such as age-specific person years data, from authors). Exploration of non-linearity should also include shape of the curve in the low exposure area.

The panelists also urged alternative approaches to meta-analyses. In particular, panelists recommended meta-regression using original (untransformed) exposure-response coefficients, in which predictor variables include the estimated percentage of amphiboles, percentage of fiber greater than 10 μm , and categorical grouping of studies according to quality. Original exposure-response coefficient variances should be used in conjunction with random effects models in which residual inter-study variation is estimated. Analyses restricted to long latency and a predictor variable for industry type should be considered. A priori distribution for inter-study residual variance might also be considered. Meta-regression will allow simple inspection of likelihoods to consider the importance of different predictor variables. Sensitivity analyses should be conducted in which the inclusion or exclusion of specific studies or groups of studies is evaluated.

Cigarette smoking. Most panelists felt strongly that future analyses need to pay more attention to the effects of smoking on the lung cancer exposure-response model and extrapolations to risk. However, the current data sets have variable and limited information available on smoking. The panelists noted that smoking is the primary cause for lung cancer, but the lung cancer dose-response relationship for smoking is complex due to the effects of smoking duration, intensity, and cessation.

The impact of smoking has effects on both the estimation and the application of the model for projecting risk of lung cancer due to asbestos exposure. This may be an especially critical issue for low-exposure extrapolation. With respect to estimation, accepting the form of the proposed model, the effect of smoking may require different K_L values for smokers and non-smokers. The panelists recognized that there is limited epidemiologic data to address this issue, but recommend that it be investigated. With respect to applying the model to make risk projections for any future cohort, the background rate of lung cancer employed in the model needs to be carefully determined to capture the smoking behavior of the cohort.

Localized tremolite exposures. During the course of public comments, the panel received input from several individuals who expressed concerns about environmental exposures to tremolite asbestos from localized geologic formations in California. The individuals suggested that inadequate attention had been given to characterization of the exposures to residents of these communities. While the panel was not in a position or charged with the evaluation of this issue, the panel did feel that this was a potentially serious matter deserving of attention by the appropriate public health authorities. Evaluation of these kinds of situations would benefit from the use of the improved risk assessment methodology being considered.

The remainder of this report summarizes the discussions and observations that led to these findings, reviews the panelists' comments on many topics not listed in this executive summary, and documents the observer comments provided at the workshop.

1. INTRODUCTION

This report summarizes a peer consultation by 11 expert panelists of a proposed protocol to assess asbestos-related risks. Contractors to the U.S. Environmental Protection Agency (EPA) developed the proposed protocol, which is documented in a report titled: “Technical Support Document for a Protocol to Assess Asbestos-Related Risk” (Berman and Crump 2001). The purpose of the peer consultation workshop was to provide EPA feedback on the scientific merit of the proposed protocol. The peer consultation workshop took place in a meeting open to the public on February 25–27, 2003, in San Francisco, California.

This report summarizes the technical discussions among the expert panelists and documents comments provided by observers. These discussions largely focused on three topic areas: interpretations of the epidemiology and toxicology literature, the proposed exposure index, and general questions about key assumptions and inferences in the protocol. The remainder of this introductory section presents background information on the protocol (Section 1.1), describes the scope of the peer consultation workshop (Section 1.2), and reviews the organization of this report (Section 1.3).

1.1 Background

EPA’s current assessment of asbestos toxicity is based primarily on an asbestos review completed in 1986 (EPA 1986) and has not changed substantially since that time. The 1986 assessment considers six mineral forms of asbestos and all asbestos fiber sizes longer than 5 micrometers (μm) to be of equal carcinogenic potency. However, since 1986, asbestos measurement techniques and the understanding of how asbestos exposure contributes to disease have improved substantially. To incorporate the knowledge gained over the last 17 years into the agency’s toxicity assessment for asbestos, EPA contracted with Aeolus, Inc., to develop a proposed methodology for conducting asbestos risk assessments. The proposed methodology distinguishes between fiber sizes and fiber types in estimating

potential health risks related to asbestos exposure. The methodology also proposes a new exposure index for estimating carcinogenic risk.

As a key step in determining the scientific merit of the proposed risk assessment methodology, EPA decided to obtain expert input on the draft report through a peer consultation workshop. The purpose of the workshop was to obtain feedback from subject-matter experts during the development stage of the proposed risk assessment methodology; the workshop was not an official peer review. Eastern Research Group, Inc. (ERG), organized and implemented the peer consultation workshop under a contract to EPA.

1.2 Scope of the Peer Consultation Workshop

The peer consultation involved many activities before the workshop (see Section 1.2.1), at the workshop (see Section 1.2.2), and after the workshop (see Section 1.2.3). The following subsections describe these activities.

1.2.1 Activities Prior to the Peer Consultation Workshop

This section describes the major activities ERG and the expert panelists conducted prior to the peer consultation workshop:

Select expert panelists. ERG selected the expert panelists for the peer consultation workshop. ERG sought to compile a panel of experts with broad experience and expertise in the following disciplines: toxicology, epidemiology, biostatistics, asbestos sampling and analytical methods, EPA's human health risk assessment guidelines, and asbestos-related environmental and occupational health issues. Appendix A lists the expert panelists ERG selected, and Appendix B includes brief biographies that summarize the panelists' areas of expertise.

Every panelist is either a senior scientist, physician, or researcher with extensive experience in the aforementioned fields, as demonstrated by peer-reviewed publications, awards, and service

to relevant professional societies. To ensure the peer consultation offered a balanced perspective, ERG intentionally selected expert panelists with a broad range of affiliations (e.g., academia, consulting, state and federal agencies). When searching for panelists, ERG asked all candidates to disclose real or perceived conflicts of interest.

- # ***Prepare a charge to the expert panelists.*** ERG worked with EPA to prepare written guidelines (commonly called a “charge”) for the peer consultation workshop. The charge includes 12 specific questions, organized into 4 topic areas. Discussions at the workshop largely addressed the technical issues raised in the charge, but the expert panelists were encouraged to discuss other relevant matters that were not specifically addressed in the charge questions. A copy of the charge is included in Appendix B.

- # ***Distribute review documents and other relevant information.*** Several weeks prior to the peer consultation workshop, ERG sent every panelist copies of the charge and the proposed risk assessment methodology (Berman and Crump 2001). These items formed the basis of the technical discussions at the workshop. In addition, ERG distributed several additional publications on related topics (see Table 1, at the end of this section, for list of the publications). The supplemental publications were provided largely in response to panelists’ requests for further background information on selected issues. The panelists also circulated publications amongst themselves on specific topics. Finally, one of the meeting chairs noted for the record that, upon arriving in San Francisco, he also received a memo and copies of many abstracts and other information from Cate Jenkins of EPA. The meeting chair offered to share these materials with other panelists during the workshop.

- # ***Obtain and compile the panelists’ premeeting comments.*** After receiving the workshop materials, the panelists were asked to prepare their initial responses to the charge questions. Booklets containing the premeeting comments were distributed to the expert panelists before the workshop and were made available to observers at the workshop. These initial comments are included in this report, without modification, as Appendix B. It should be noted that the premeeting comments are preliminary in nature. Some panelists’ technical findings may have changed after the premeeting comments were submitted.

1.2.2 Activities at the Peer Consultation Workshop

The 11 expert panelists and approximately 75 observers attended the peer consultation workshop, which was held at the Westin St. Francis Hotel in San Francisco, California, on February 25–27, 2003. The workshop was open to the public, and the workshop dates and times were announced in the

Federal Register. Appendix C lists the observers who confirmed their attendance at the workshop registration desk. The workshop schedule generally followed the agenda, presented here as Appendix D.

The workshop began with introductory remarks from Ms. Jan Connery (ERG), the facilitator of the peer consultation. Ms. Connery welcomed the expert panelists and observers, stated the purpose of the workshop, identified the document being reviewed, and explained the procedure for observers to make comments. Mr. Richard Troast (EPA) then provided background information on the review document and EPA's ongoing efforts to assess asbestos toxicity (see Section 1.1). Mr. Troast identified the main differences between EPA's existing asbestos risk assessment methodology (EPA 1986) and the proposed methodology (Berman and Crump 2001). Mr. Troast noted that the expert panelists' feedback will ultimately help EPA complete its update of asbestos health risks for the Integrated Risk Information System (IRIS); he clarified that the final IRIS update will be subject to peer review or Science Advisory Board review before being implemented. Following these opening remarks, Dr. Wayne Berman and Dr. Kenny Crump—the authors of the proposed methodology—presented detailed information on the review document; Section 2 of this report summarizes their presentations.

After the background presentation, Dr. Roger McClellan and Dr. Leslie Stayner chaired the technical discussions that followed. For the remainder of the meeting, the panelists engaged in free-flowing discussions when answering the charge questions and addressing additional topics not specified in the charge. Observers were given the opportunity to provide verbal comments three different times during the workshop; these observer comments are documented in Appendix E. Representatives from EPA and the document authors provided clarifications on the proposed methodology periodically throughout the 2½-day workshop.

1.2.3 Activities Following the Peer Consultation Workshop

The primary activity following the peer consultation workshop was preparing this summary report. A technical writer from ERG who attended the meeting prepared a draft of this report, which ERG distributed to the 11 expert panelists and asked them to verify that the draft accurately reflects the tone and substance of the panelists' discussions at the workshop. After incorporating the panelists' suggested revisions to the draft report, ERG submitted the final report (i.e., this report) to EPA.

1.3 Report Organization

The structure of this report follows the order of the technical discussions during the meeting. Section 2 summarizes Dr. Berman and Crump's background presentations. Sections 3 through 6 are records of the panelists' discussions on the four main topic areas: interpretations of the epidemiology and toxicology literature (Section 3), the proposed exposure index (Section 4), general questions (Section 5), and conclusions and recommendations (Section 6). Finally, Section 7 provides references for all documents cited in the text.

The appendices to this report include background information on the peer consultation workshop. This information includes items that were on display at the workshop and items generated since the workshop (e.g., a final list of attendees). The appendices contain the following information:

- # List of the expert panelists (Appendix A).
- # The panelists' premeeting comments, the charge to the reviewers, and brief bios of the expert panelists (Appendix B).
- # List of registered observers of the peer consultation workshop (Appendix C).
- # Agenda for the peer consultation workshop (Appendix D).
- # Observer comments provided at the peer consultation workshop (Appendix E).
- # Observer post-meeting comments (Appendix F).

Table 1
References ERG Provided to the Expert Panelists

Berman, DW and Crump K. 1999. Methodology for Conducting Risk Assessments at Asbestos Superfund Sites; Part 1: Protocol. Final Draft. Prepared for U.S. Environmental Protection Agency. February 15, 1999.
Berman, DW and Crump K. 2001. Technical Support Document for a Protocol to Assess Asbestos-Related Risk. Final Draft. Prepared for U.S. Department of Transportation and U.S. Environmental Protection Agency. September 4, 2001.
Berman, DW, Crump, K., Chatfield, E., Davis, J. and A. Jones. 1995. The Sizes, Shapes, and Mineralogy of Asbestos Structures that Induce Lung Tumors or Mesothelioma in AF/HAN Rats Following Inhalation. Risk Analysis. 15:2,181-195.
Berman, DW. 1995. Errata. Risk Analysis. 15:4, 541.
Committee on Nonoccupational Health Risks of Asbestiform Fibers. Breslow, L., Chairman. 1984. Asbestiform Fibers Nonoccupational Health Risks. Washington, DC: National Academy Press.
EPA 1986. Airborne Asbestos Health Assessment Update. U.S. Environmental Protection Agency. EPA 600/8-84-003F. 1986.
NIOSH Interdivisional Fiber Subcommittee Report. Prepared by the NIOSH Interdivisional Fiber Subcommittee. 1999.

2. BACKGROUND ON THE PROPOSED PROTOCOL TO ASSESS ASBESTOS-RELATED RISK

This section summarizes presentations given by the principal authors of the proposed risk assessment methodology. These presentations were given because several panelists asked ERG, prior to the peer consultation workshop, if the authors would provide detailed background information on how the methodology was developed. This section reviews the major presentation topics, but does not present the panelists' comments on the proposed protocol. Sections 3 through 6 document the expert panelists' technical feedback on the protocol.

Motivation for developing the proposed protocol. Dr. Berman identified several reasons for developing the updated protocol for assessing asbestos-related risks. These reasons include EPA's existing asbestos models being inconsistent with inferences from the scientific literature, the need for having uniformly-applied sampling and analytical procedures to measure asbestos characteristics most predictive of risk, and the belief that EPA's current asbestos risk assessment methodology may not be adequately protective in some circumstances. To improve upon the current methodology, the authors intended to develop a risk assessment model that adequately predicts cancer risk in all studied environments and can therefore be applied with much greater confidence to environments that have not been studied. Dr. Berman outlined the general approach taken to develop the proposed protocol, as summarized in the following bulleted items.

Dr. Berman provided background information on and definitions for asbestos, other fibrous structures, asbestos morphology, and cleavage fragments. He also described the capabilities and limitations of the analytical techniques that have been used to characterize asbestos exposures, such as midget impingers, phase contrast microscopy (PCM), scanning electron microscopy (SEM), and transmission electron microscopy (TEM). Dr. Berman explained how differences in these analytical techniques must be critically evaluated when comparing results reported in all epidemiological and other types of studies that examine asbestos exposure. Dr. Berman also stressed that it is not just differences in analytical techniques, but choice of specific methods for each analytical technique that affects results. Further information on these topics is included in Chapter 4 of the proposed protocol (Berman and Crump 2001).

Re-analysis of human epidemiological data. Dr. Crump described how the authors evaluated the human epidemiological data. He displayed a list of the studies that were considered, noting that he had access to raw, individual-level data for three occupational cohorts: chrysotile textile workers in South Carolina, United States; crocidolite miners in

Wittenoom, Australia; and chrysotile miners and millers in Quebec, Canada. All data sets with exposure data were considered in the analysis, and criteria were not established for selecting studies. Dr. Crump then presented findings for asbestos-related risks for lung cancer and mesothelioma.

For lung cancer, Dr. Crump first reviewed EPA's existing lung cancer model for asbestos exposure (see equation 6.1 in the proposed protocol), which relates the relative risk of lung cancer mortality linearly to cumulative asbestos exposure, with a 10-year lag time. Dr. Crump noted that the model predicts that relative risk for developing lung cancer remains constant after asbestos exposure ceases—an assumption he showed was reasonably consistent with findings from epidemiological studies. Dr. Crump also discussed how the model assesses interactions between exposures to cigarette smoke and to asbestos—an issue the panelists revisited several times later in the workshop (e.g., see Section 3.1.1 and the executive summary). Dr. Crump presented a series of tables and figures demonstrating the adequacy of multiple lung cancer models: first using EPA's existing lung cancer model, next using a modified version of the model that accounts for differences in the background rates of lung cancer, and finally using the proposed lung cancer model, which considers an exposure index that assigns greater carcinogenic potency to amphibole fibers and to longer fibers.

Similarly, Dr. Crump reviewed the performance of EPA's mesothelioma model for asbestos exposures (see equation 6.11 in the proposed protocol), which predicts that mesothelioma risks vary linearly with the average asbestos exposure and increase quadratically with time from onset of exposure. Dr. Crump presented several tables and graphs indicating how well EPA's existing model and the proposed protocol fit the human epidemiological data. He made several conclusions about the existing risk model, including that mesothelioma risk coefficients varied considerably across the cohorts and the risk coefficients were generally higher for cohorts exposed primarily to amphibole fibers, compared to those exposed primarily to chrysotile fibers. Dr. Crump also noted that the data did not support consideration of a sub-linear or threshold dose-response relationship. This latter point generated considerable discussion later in the workshop (e.g., see Section 4.3).

Dr. Crump then described the meta-analysis the authors conducted to evaluate the relative potency of amphibole and chrysotile fibers. First, he explained how the authors weighted the different studies in the meta-analysis, based on uncertainty factors assigned to the individual studies. Dr. Crump identified the four uncertainty factors and described generally how each factor was assigned. Sources of uncertainty included representativeness of air sampling data, the availability of conversion factors to express exposures in terms of PCM concentrations, and whether data on exposure duration were available. Dr. Crump then highlighted the main conclusions from the meta-analysis. For lung cancer, the meta-analysis suggested that amphibole fibers are approximately five times more potent than are chrysotile fibers, but the difference in potency was not statistically significant (i.e., the authors could not reject the hypothesis that

chrysotile fibers and amphibole fibers are equally potent). For mesothelioma, the meta-analysis suggested that chrysotile fibers are 0.002 times as potent as amphibole fibers, and the difference in potency was statistically significant.

- # ***Inferences drawn from the broader literature.*** Dr. Berman described how the authors incorporated inferences from the broader scientific literature into the proposed protocol. He reviewed key findings on how various mechanisms are biologically related to how asbestos causes disease. These mechanisms included respiration, deposition, degradation, clearance, translocation, and tissue-specific biological responses. Chapter 7 of the review document provides detailed information on the relevance of these mechanisms, with emphasis on the influence of fiber type and fiber dimension.

- # ***Derivation of the exposure index.*** Dr. Berman explained how the authors derived the exposure index, which is largely based on an earlier re-analysis (Berman et al. 1995) of six animal inhalation studies conducted by a single laboratory. That re-analysis found that lung tumor incidence is adequately predicted using an exposure index that assigns no carcinogenic potency to fibers shorter than 5 μm , relatively low carcinogenic potency to fibers with lengths between 5 and 40 μm and diameters less than 0.4 μm , and the greatest carcinogenic potency to fibers longer than 40 μm and thinner than 0.4 μm . However, these findings could not be applied directly to the human epidemiological data, because the epidemiological studies do not include exposure measurements that quantify the relative amounts of asbestos fibers shorter and longer than 40 μm .

Dr. Berman noted that the proposed protocol includes an *ad hoc* assumption that the fiber size weighting factors optimized from the laboratory animal studies can be applied to humans, but with a length cut-off of 10 μm in the exposure index, rather than a cut-off of 40 μm . Dr. Berman emphasized that this assumption was made to model the critical characteristics of asbestos in a manner that reasonably captures cancer risks observed across multiple epidemiological studies. He acknowledged that asbestos potency is likely a continuous function of fiber length, but the exposure measurements from the available animal and epidemiological studies do not support incorporating such a continuous function in the exposure-response model. The panelists commented on the proposed exposure index when discussing topic area 3 (see Section 4).

Dr. Berman also noted that the authors selected a conservative set of dose-response coefficients (see Table 6-30 of the review document), rather than using the optimized ones from the animal studies (see Table 6-29). However, the conservative and optimized dose-response coefficients were reasonably consistent: none of the conservative coefficients differed by more than a factor of 4 from the corresponding optimized ones.

- # ***Conclusions regarding proposed protocol.*** Dr. Berman indicated that the proposed protocol is substantially more consistent with inferences documented in the scientific literature (i.e., that

long, thin structures contribute most to risk) than EPA's existing risk assessment methodology. Further, the proposed protocol provides a better fit to cancer risks observed in the human epidemiological studies than does EPA's existing model, and the proposed protocol appears to underestimate risks of lung cancer and mesothelioma less frequently and to a lesser degree than the existing approach. Finally, by recommending use of a standardized analytical method that links directly to the exposure index, the proposed protocol will help ensure that future risk assessments are conducted in a consistent fashion and their results can be readily compared from one study to the next.

3. COMMENTS ON TOPIC AREA 1: INTERPRETATIONS OF THE EPIDEMIOLOGY AND TOXICOLOGY LITERATURE

This section summarizes the panelists' discussions on the interpretations of the epidemiology and toxicology literature. The meeting co-chairs—Dr. McClellan and Dr. Stayner—facilitated the discussions on this topic area, which focused first on lung cancer (see Section 3.1) and then on mesothelioma (see Section 3.2). This section presents a record of discussion of the topics mentioned during the workshop. Several panelists referred to their premeeting comments (see Appendix B) for additional suggestions for how the review of epidemiology and toxicology literature can be improved.

3.1 Lung Cancer

The panelists discussed at length whether the epidemiology and toxicology literature support the proposed protocol's finding for how lung cancer potency varies with fiber type and fiber length. This section summarizes these discussions, first on fiber type (Sections 3.1.1 and 3.1.2) and then on fiber length (Sections 3.1.3 and 3.1.4). General issues regarding the lung cancer evaluation are presented in Section 3.1.5.

3.1.1 Lung Cancer and Fiber Type: Inferences from the Epidemiology Literature

According to the proposed risk assessment methodology, amphibole fibers have a 5-fold greater lung cancer potency than do chrysotile fibers. The panelists had differing opinions on whether this finding is consistent with the epidemiology literature. On the one hand, some panelists indicated that the epidemiology literature is consistent with amphibole fibers being more potent for lung cancer, though the magnitude of this increase may not be known precisely. One panelist noted, for example, that multiple analyses (e.g., Hodgson and Darnton 2000, Berman and Crump 2001, and the statistical analyses a panelist presented during this discussion) all point to a consistent increased lung cancer potency for amphibole fibers compared to chrysotile fibers, albeit a small increase. On the other hand, other

panelists did not believe the epidemiology literature supports this conclusion, for reasons stated below. Finally, other panelists were not convinced that the epidemiology literature supports the higher lung cancer potency for amphibole fibers, but they believed the difference in potency seems likely based on evidence from the animal toxicology studies (see Section 3.1.3) and lung burden studies. A summary of the panelists' discussion on this topic follows:

Comments on specific publications. Several panelists cited specific studies to support their positions on the relative lung cancer potency of chrysotile and amphibole fibers, but the panelists often had differing opinions on the inferences that should be drawn. The panelists mentioned the following specific studies:

- ▶ Some panelists noted that a recent re-analysis of 17 cohorts (Hodgson and Darnton 2000) indicates that the lung cancer potency for amphibole fibers is 10 to 50 times greater than that for chrysotile fibers. One panelist did not agree with this finding, due to the crude approach the article uses to characterize relative potency. Specifically, this panelist noted that carcinogenic potency was calculated by dividing the overall relative risk for a given cohort by the average exposure for the entire cohort, even for cohorts where the data support more sophisticated exposure-response modeling. He was particularly concerned about the authors' decision to omit the cohort of South Carolina textile workers from the meta-analysis. This decision was apparently based on the South Carolina cohort being an outlier, due to its much higher lung cancer potency when compared to other studies. The panelist noted, however, that the lung cancer risk for the South Carolina cohort is not unusually high when compared to other cohorts of textile workers. The panelist was concerned that omitting this study might have biased the article's finding regarding relative lung cancer potency. No other panelists discussed the review article.
- ▶ One panelist cited a study of Quebec chrysotile miners and millers (Liddell et al. 1997, 1998) that reports that increased lung cancer risk was limited to the mining region with the highest level of tremolite asbestos, after correction for smoking and exposure. The article was distributed to the panelists on the first day of the workshop, but no panelists commented further on the study.
- ▶ One panelist noted that his review of multiple textile cohorts (Stayner, Dankovic, and Lemen 1996) found relatively small differences in lung cancer potency, even though some of the cohorts were exposed to asbestos mixtures containing different proportions of amphibole fibers.

- ▶ One panelist indicated that further evidence on how fiber types relates to lung cancer potency can be gleaned from epidemiological studies that were not included in the meta-analysis due to inadequate exposure data for exposure-response modeling. Examples include a study of non-occupationally exposed women from two chrysotile asbestos mining regions (Camus et al. 1998) and a study of railroad workers employed by shops that processed different proportions of amphibole fibers (Ohlson et al. 1984). Both studies, she noted, provide evidence that amphibole fibers exhibit greater lung cancer potency. This panelist added that studies of auto mechanics have provided no convincing evidence of increased lung cancer due to chrysotile exposure, though she acknowledged that the absence of an effect might reflect the short fiber length in the friction brake products. One panelist cautioned about inferring too much from these studies regarding fiber type because they were not controlled for other factors, such as fiber length and level of exposure.

- ▶ One panelist added that a recent study of a cohort of Chinese asbestos plant workers (Yano et al. 2001) should be considered in future updates to the proposed protocol; the workers in the cohort had increased risks for lung cancer and were reportedly exposed to “amphibole-free” chrysotile asbestos. However, another panelist cited a publication (Tossavainen et al. 2001) that indicates that asbestos from many Chinese chrysotile mines actually does contain varying amounts of amphibole fibers.

- ▶ Several panelists noted that the proposed protocol’s meta-analysis found a 5-fold difference in lung cancer potency between amphibole and chrysotile fibers. However, other panelists indicated that the reported difference was not statistically significant. Some panelists had additional reservations about the authors’ meta-analysis, as summarized in the following bulleted items.

Comments on the meta-analysis approach. Several panelists commented on alternate approaches the authors could have used to conduct their meta-analysis of the epidemiology studies. One panelist noted that the lung cancer potencies reported by the various studies exhibit considerable heterogeneity. In such cases, meta-regression is conventionally used to identify which factors account for the variability in the results (i.e., in the lung cancer potencies). This panelist suggested that the meta-analysis should have considered other factors in addition to fiber type and dimension; such other factors could include industry, follow-up time for the cohort, and estimated percentage of amphibole fibers in the exposures, to the extent that data on these other factors are available.

To demonstrate how more detailed investigation might reveal further insights, one panelist presented his own initial statistical analysis of the epidemiological studies. This analysis used a fixed effects model and a random effects model, both inverse weighted by the variance of the studies. His analysis examined how industry and fiber type contribute to the heterogeneity

observed among the cohorts and found that the industry of the cohort appears to be a stronger predictor than fiber type. The panelist explained that the purpose of displaying his statistical analysis was to highlight how other approaches to conducting meta-analysis can offer different insights on the epidemiological data. This panelist recommended that the authors conduct similar meta-regression analyses to investigate the importance of various variables on the lung cancer potency.

This panelist also demonstrated how a sensitivity analysis might yield additional information on influential studies. Using a fixed effects model, the panelist first showed how lung cancer potency factors (K_L) vary with exposure to chrysotile fibers, amphibole fibers, and mixed fiber types. When all epidemiological studies were considered in his analysis, the amphibole fibers were found to be three times more potent than the chrysotile fibers. When the cohort of chrysotile miners and millers from Quebec was omitted from this analysis, however, the amphibole fibers were found to be nearly two times *less* potent than the chrysotile fibers. Conversely, when the cohort of textile workers from South Carolina was omitted, the amphibole fibers were found to be more than ten times more potent than the chrysotile fibers. Given that the conclusions drawn about the relative potency of chrysotile and amphibole fibers appear to be highly sensitive to whether single studies are omitted from the analysis, this panelist was more skeptical about whether the increased potency of amphibole fibers is a robust finding. He recommended that the authors, when completing the proposed protocol, conduct similar sensitivity analyses to help reveal the factors or studies that appear to contribute most to lung cancer.

Another panelist agreed with this feedback, and provided further comments on the meta-analysis, noting that these analyses typically start with establishing criteria for study inclusion. After selecting studies to evaluate, she said, various statistical analyses can be used to test hypotheses and to understand the concordance and disparity among the individual studies. The panelist thought such an approach is needed to help understand the variability in potency factors observed across the multiple studies and to identify for further analysis the studies found to be most descriptive of exposure-response. To clarify the authors' approach, Dr. Berman indicated that the meta-analysis considered any published epidemiological study with sufficient quantitative exposure data that allowed for a reasonable estimate of the exposure-response relationship; uncertainty factors were then assigned to give greatest weight to the most robust studies. In response, additional panelists concurred with the original comment that meta-analyses conventionally begin with establishing explicit study inclusion criteria. These panelists clarified that they are not advocating removing a majority of studies currently considered in the proposed protocol, but rather being more judicious in selecting the studies to evaluate.

One panelist offered additional comments on the meta-analysis. He supported, for instance, the use of sensitivity analyses, and encouraged the authors to conduct additional analyses to identify influential studies, factors that contribute to risk, and the impact of different weighting factors. The panelist also noted that more sophisticated statistical methodologies (e.g., Bayesian

modeling, Markov Monte Carlo) can be used to generate distributions of outputs, rather than discrete values, which might offer greater understanding of the inferences that can be drawn from the epidemiological studies.

Disparate findings from the South Carolina and Quebec cohorts. Multiple panelists noted that the issue of the relative lung cancer potency of chrysotile and amphibole fibers depends largely on how one interprets the disparate findings from the cohort of textile workers in South Carolina and the cohort of chrysotile miners and millers in Quebec. Two of these panelists indicated that the relative potency issue likely will not be resolved until the underlying reasons for the differences between these two studies are better understood. The other panelist viewed the difference in potency observed across industries (i.e., mining versus textile) as a more important matter than the difference between the two specific cohorts. When discussing these studies, two panelists indicated that the increased lung cancer risk for the South Carolina cohort might be attributed to exposure to amphibole fibers, which are known to be found in trace levels in commercial chrysotile.

Relevance of fiber durability. One panelist noted that the issue of fiber durability often enters the debate on the relative lung cancer potency of chrysotile and amphibole fibers. Though he agreed that the animal toxicology data indicate that amphibole fibers are more persistent than chrysotile fibers, the panelist noted that trends among the human epidemiological data—particularly the fact that lung cancer risk does not appear to decrease with time since last exposure, even for chrysotile—suggest that the lower durability of the chrysotile fibers might not be important.

Influence of smoking. The panelists had differing opinions on how the proposed protocol should address cigarette smoking. In terms of inferences drawn from the epidemiological literature, two panelists noted that very limited data are available on smoking, making quantitative analysis of its interactions with asbestos exposures difficult. Specifically, only one study includes detailed information on smoking, but that study found no difference in lung cancer potency between smokers and non-smokers. During this discussion, Dr. Berman explained that the proposed protocol assumes a multiplicative interaction between smoking and asbestos exposure, consistent with EPA's 1986 model. Dr. Berman noted that a multiplicative factor in the model, α , represents the background risk in the studied cohort relative to the risk in the comparison population, and both groups include smokers; he added that the influence of smoking is addressed implicitly in the model because it is a relative risk model in which the effect of asbestos is multiplied to the background risk that is present. A panelist clarified, however, that neither the potency factors nor α were derived based on observations of smoking prevalence in the epidemiological studies.

One panelist emphasized that the confounding effects of smoking greatly complicates the analysis of lung cancer potency. He noted that the relative lung cancer risk from asbestos exposure is

considerably lower than that for cigarette smoking. As a result, the panelist wondered how the meta-analysis can truly discern the relative potency of the asbestos fiber types from studies that present no information on cigarette smoking. This panelist provided an example to illustrate his concern: if a given cohort has between 5 and 10% more smokers than the typical population, this increased prevalence of smoking alone could totally confound relative risks attributed to asbestos. The panelist indicated that all future analyses of epidemiological data will suffer from similar limitations, so long as detailed information on smoking is not available.

General comments. During this discussion, some panelists offered several general comments that apply to the entire proposed protocol. These comments included concerns about the transparency of the analyses, questions about data tables being inconsistent with text in the body of the report, and some panelists' inability to reproduce certain findings from the available data. These general comments are reflected in the executive summary of this report.

3.1.2 Lung Cancer and Fiber Type: Inferences from Animal Toxicology and Mechanistic Studies

The panelists offered varying insights on the inferences that can, or should, be drawn from animal toxicology studies and mechanistic studies regarding the relative lung cancer potency for chrysotile and amphibole fibers.

Citing various publications (e.g., Lippmann 1994), multiple panelists noted that the animal toxicology studies do not support the 5-fold difference in lung cancer potency between chrysotile and amphibole fibers. Two panelists added that the absence of different potencies might result from the animal studies being of too short duration (typically no longer than 2 years) for the greater dissolution of chrysotile fibers to be an important factor. Another panelist added that exposure levels in some animal studies are not relevant to human exposures; as an example, he noted that a recent rat inhalation study (Hesterberg et al. 1998) involved exposure levels at 11,000 fibers per cubic centimeter. These panelists indicated that the animal studies are generally more informative of how lung cancer potency varies with fiber length (see Section 3.1.4), and are less informative on how potency varies with fiber type.

The panelists noted that *in vitro* studies exhibit various findings, depending on the study design and endpoint assessed. One panelist, for instance, indicated that some *in vitro* studies suggest that chrysotile fibers are actually more potent than amphibole fibers. Other panelists added that many *in vitro* studies show crocidolite being considerably more toxic than chrysotile. These panelist cautioned against drawing firm conclusions from the *in vitro* studies, however, given that the study duration is far too short for any impact of dissolution to be observed. Finally, another panelist referred to the International Agency for Research on Cancer (IARC) consensus statement on fiber carcinogenesis for an overview of inferences that can be drawn from mechanistic studies: “Overall, the available evidence in favor of or against any of these mechanisms leading to the development of lung cancer and mesothelioma in either animals or humans is evaluated as weak” (IARC 1996).

Based on the previous comments, the panelists cautioned about attempting to draw inferences from the animal toxicology for several reasons. One panelist indicated that the animal studies have limited utility because lung cancer in humans results from a complex set of exposures, including cigarette smoke, and because rats, when compared to humans, develop different types of tumors at different sites. Another panelist reiterated that the duration of most animal studies precludes one from observing dissolution effects. Given these limitations, two panelists emphasized that conclusions should be based primarily on the epidemiological data, especially considering the volume of human data that are available. Though not disagreeing with this recommendation, one panelist noted that the exposure index—one of the major outcomes of the proposed protocol—is, in fact, based on observations from animal studies.

3.1.3 Lung Cancer and Fiber Dimension: Inferences from the Epidemiology Literature

The panelists made several observations regarding what can be inferred from the epidemiology literature on how lung cancer potency varies with fiber dimension, though they first noted that most published epidemiology studies do not include detailed data on the distribution of fiber dimensions to which cohorts were exposed. Overall, the panelists generally agreed that indirect evidence from the

epidemiological studies supports the proposed protocol's finding that longer fibers have greater carcinogenic potency for lung cancer. They added, however, that the epidemiology literature provides no evidence to support or refute the magnitude of the relative potencies used in the proposed protocol (i.e., fibers longer than 10 μm being 300 times more potent than those with lengths between 5 and 10 μm). The panelists made no comments about fiber diameter when discussing this matter. Specific discussion topics follow:

- # ***Observations from the epidemiology literature.*** The panelists identified several studies that provide general insights on the role of fiber size in lung cancer. One panelist, for instance, noted that cohorts of textile workers, which were believed to be exposed to relatively longer asbestos fibers, exhibit higher lung cancer relative risks than do cohorts of miners or cement product workers. Another panelist indicated that studies of taconite miners from Minnesota (Cooper et al. 1988) and gold miners from South Dakota (McDonald et al. 1978) found no increased lung cancer risks among the cohorts, which were known to be exposed primarily to fibers shorter than 5 μm (see Dr. Case's premeeting comments for further information on these studies). This panelist added that the Minnesota Department of Health is currently updating the study on taconite miners and a publication is pending. Another panelist added that epidemiology studies of workers exposed to asbestos from friction brake products show no clear evidence of increased lung cancer. This panelist acknowledged that these epidemiology studies do not include exposure measurements, but other studies of this work environment have indicated that the asbestos fibers in friction brake products are predominantly short chrysotile fibers.

- # ***Relevance of fibrous structures shorter than 5 μm .*** Some panelists noted that no epidemiology studies have examined the relative potency specifically of fibrous structures shorter than 5 μm , thus no conclusions could be drawn from the epidemiology studies alone. While not disagreeing with this observation, one panelist reminded panelists that airborne particles and fibers have a broad distribution of fiber lengths, with a clear majority (75–90%) of fibrous structures being shorter than 5 μm . This panelist added that indirect inferences can be drawn from the epidemiology studies listed in the previous bulleted item. Another panelist noted that the fibrous structures shorter than 5 μm behave more like particles rather than fibers, at least in terms of lung deposition and clearance patterns. Finally, two panelists indicated that an ATSDR expert panel recently evaluated the issue of relative potency of fibers shorter than 5 μm ; however, the final report from that expert panel meeting was not available until after the peer consultation workshop. The final report has since been released, and a conclusion from that panel was that “there is a strong weight of evidence that asbestos and synthetic vitreous fibers shorter than 5 μm are unlikely to cause cancer in humans” (ERG 2003).

- # ***Statistical analyses in the proposed protocol.*** As indirect evidence that longer fibers have greater carcinogenic potency, one panelist indicated that the exposure-response modeling by Drs. Berman and Crump showed an improved fit to the observed relative risk from epidemiology studies when using an exposure index that assigns greater weight to longer fibers and no risk to fibers shorter than 5 μm . Another panelist concurred, but added that the authors could have attempted to determine the specific weighting (i.e., between longer and shorter fibers) that would optimize the fit to the epidemiological studies.

3.1.4 Lung Cancer and Fiber Dimension: Inferences from Animal Toxicology and Mechanistic Studies

The panelists generally agreed that the animal toxicology studies and mechanistic studies indicate that fiber dimension—especially fiber length—plays an important role, both in terms of dosimetry and pathogenesis. However, panelists had differing opinions on the specific cut-offs that should be used for fiber diameters and lengths in the exposure-response modeling (though panelists generally concurred that fibers shorter than 5 μm should be assigned zero potency).

- # ***Fiber length.*** Multiple panelists noted that the animal toxicology studies provide compelling evidence that lung cancer potency increases with fiber length. Another panelist agreed, but had reservations about assigning no potency to fibrous structures shorter than 5 μm , based on a recent study of refractory ceramic fibers (Bellman et al. 2001) that found that the incidence of inflammation and fibrosis appears to be related to the presence of small fibers in the lung. This panelist indicated that exposure to small fibers likely has some bearing on the oxidative stress state and inflammation in the lung, and he suspected that the exposure-response relationship for long fibers might depend on co-exposures or past exposures to shorter fibers. Based on these observations, the panelist was hesitant to exclude fibrous structures shorter than 5 μm from the proposed risk assessment methodology. On the other hand, another panelist added that animal toxicology studies have shown that fibrosis endpoints are strongly related to fiber length, with exposures to shorter fibers showing less evidence of fibrosis or lung damage. The panelists revisited the significance of fibers shorter than 5 μm when discussing the proposed exposure index (see Section 4).
- # ***Fiber diameter.*** The panelists offered several comments on the role of fiber diameter in the proposed protocol. Noting that fibers with diameters up to 1.5 μm are capable of penetrating to sensitive portions of the lung during oral inhalation, one panelist indicated that this range of fiber diameters should not be excluded from future risk assessments. Other panelists shared the

concern of assigning no lung cancer potency to respirable fibers with diameters greater than 0.5 μm , especially considering that respirability patterns in laboratory animals differ from those in humans (i.e., thicker fibers are more likely to deposit in the human lung than they are in the rat lung).

The panelists also discussed a statement in the proposed protocol that “few fibers thicker than 0.7 μm appear to reach the deep lung.” First, one panelist indicated that the proposed protocol includes outdated information on fiber deposition patterns; he recommended that the authors obtain more current insights from specific publications (e.g., Lippmann 1994) and from the latest lung dosimetry model developed by the International Commission on Radiological Protection. Second, another panelist questioned the relevance of deposition in the deep lung, because humans tend to develop bronchogenic carcinomas, while rats develop bronchoalveolar carcinomas. Another panelist cautioned against inferring that asbestos fibers must deposit on bronchial airways to cause lung cancer in humans, noting that significant accumulation of asbestos fibers does not occur in the airways where carcinomas develop in humans, due primarily to mucociliary clearance; this panelist suspected that deposition of fibers in the deep lung is likely related to lung cancer formation in humans, though the mechanisms of carcinogenesis are not fully understood.

3.1.5 Other Issues Related to Lung Cancer

The panelists discussed several additional issues related to the proposed protocol’s evaluation of lung cancer potency. Most of the discussion focused on the utility of non-linear exposure-response modeling, but other topics were also addressed:

- # ***Consideration of non-linear exposure-response models.*** The panelists had differing opinions on the extent to which the proposed protocol should consider non-linear exposure-response modeling. On the one hand, one panelist strongly recommended that EPA consider exploring the applicability of non-linear exposure-response models, given his concerns with linear low-exposure extrapolation. This panelist acknowledged that the revised linear model in the proposed protocol clearly provides an improved statistical fit to the epidemiological data when compared to EPA’s 1986 lung cancer model, but he advocated more detailed exploration of non-linear cancer risk models, particularly to account for observations of cohorts with low exposures. This panelist was particularly concerned about the cancer risks that would be predicted for low exposures: because the slope in any linear lung cancer model will be determined largely by highly-exposed individuals, he questioned whether the slope derived from

high exposures truly applies to lowly-exposed individuals. To demonstrate his concern, this panelist indicated that the epidemiological studies consistently show that cohorts (or subsets of cohorts) with low exposure generally exhibit no increased lung cancer risk (standardized mortality ratios not statistically different from 1.0). To account for the possibility of a threshold or non-linearity in the exposure-response relationship, this panelist recommended that EPA investigate alternate exposure-response models, such as linear-linear models (i.e., models with two linear exposure-response regions having different slopes) or log-linear models.

Other panelists generally supported these comments. One panelist, for instance, noted that EPA's Draft Revised Guidelines for Carcinogen Risk Assessment indicates that exposure-response relationships should first be evaluated over the range of exposure observations, and then various approaches to extrapolate to exposure levels outside (i.e., below) this range should be investigated. Another panelist added that some studies finding no evidence of lung cancer risks among large cohorts with low exposures should factor into the decision of whether the lung cancer model should include thresholds; he cited a study of non-occupationally exposed women from chrysotile mining regions in Canada (Camus et al. 1998) to illustrate his concern. Other panelists noted that the utility of this study is limited, because exposures were not measured for individuals; further, a panelist clarified that approximately 5% of the individuals considered in this study were occupationally exposed. Finally, one panelist indicated that evidence from the epidemiology literature strongly suggests there are asbestos exposure levels below which lung cancer will not occur; this panelist added that he is unaware of any epidemiological study that has found evidence of lung cancer risk at exposure levels below 25 fiber-years. He recommended that the proposed protocol at least acknowledge the lowest exposure level at which lung cancer effects have been demonstrated.

On the other hand, some panelists were not convinced of the utility of conducting detailed analyses at low exposures and investigating possible thresholds. One panelist, for instance, indicated that a meaningful quantitative analysis of potential thresholds will not be possible, so long as the authors do not have access to raw data from additional epidemiological studies. Further, this panelist suspected that the protocol authors would find considerable heterogeneity among exposure-response slopes for low exposures, and he questioned what conclusions could be drawn by focusing exclusively on the low exposure region. Another panelist agreed, adding that the failure to find significantly increased cancer risks among lowly-exposed cohorts very likely results from poor statistical power and other uncertainties, and not necessarily from the presence of an actual exposure threshold for asbestos-related lung cancer. Finally, one panelist indicated that the National Institute for Occupational Safety and Health (NIOSH) previously examined a threshold model for the cohort of South Carolina textile workers, and that analysis revealed that the best fit of the exposure-response data was a threshold of zero (i.e., the best fit indicated that there was no threshold).

- # ***Consideration of cigarette smoking.*** Several times during the workshop, the panelists debated the ability of the proposed risk assessment model to address interactions between cigarette smoking and asbestos exposure. One panelist recommended that the authors review a recent study that examined the role of cigarette smoking on lung cancer among chrysotile miners and millers in Quebec, Canada (Liddell and Armstrong 2002). Although the panelists generally agreed that smoking is an important consideration for developing and applying the model, some panelists were not convinced that the available data are sufficient to develop an exposure-response model that accurately portrays the interactive effects of asbestos exposure and smoking. The panelists further discussed this issue further later in the workshop.

- # ***Transparency of the proposed protocol.*** Several panelists indicated that the review of epidemiological data in the proposed protocol is not presented in a transparent fashion. One panelist, for instance, sought more information on the uncertainty factors used in the meta-analysis, such as what ranges of factors were considered, what criteria were used to assign the factors, and a table of the factors that were eventually applied. This panelist also recommended that the proposed protocol identify the α -values that were determined for each epidemiological study and provide explanations for any cases when these values are unexpectedly large. Another panelist indicated that the proposed protocol should more clearly differentiate conclusions that are based on a meta-analysis of many epidemiological studies from conclusions that are based on a detailed review of just one or two studies.

- # ***The need to obtain additional raw data sets.*** The panelists unanimously agreed that EPA should make every effort to try to obtain additional raw data sets for the epidemiology studies, such that the authors can further test how adequately the proposed risk assessment model predicts risk. The executive summary of this report presents the panelists' specific recommendation on this issue.

3.2 Mesothelioma

The following paragraphs document the panelists' responses to charge questions regarding inferences from the epidemiology and toxicology literature on how mesothelioma potency varies with fiber type (Sections 3.2.1 and 3.2.2) and fiber length (3.2.3 and 3.2.4).

3.2.1 Mesothelioma and Fiber Type: Inferences from the Epidemiology Literature

The expert panelists unanimously agreed that the epidemiology literature provides compelling evidence that amphibole fibers have far greater mesothelioma potency than do chrysotile fibers—a finding reported both in the review document (Berman and Crump 2001) and a recent re-analysis of 17 cohort studies (Hodgson and Darnton 2000) that reported at least a 500-fold difference in potency. Two panelists commented further that the epidemiology literature provides no scientific support for chrysotile exposures having a role in causation of mesothelioma—an observation that is generally consistent with the meta-analysis in the proposed protocol, which failed to reject the hypothesis that chrysotile fibers have zero potency for mesothelioma.

The most notable response to this charge question was the agreement among most panelists that amphibole fibers are at least 500 times more potent than chrysotile fibers for mesothelioma, as supported by two separate reviews of epidemiological studies. The panelists made additional comments on specific matters when responding to this question, as summarized below, but the key point in this discussion was the agreement that chrysotile is a far less important cause of mesothelioma than are amphiboles.

Relative roles of chrysotile and amphibole. One panelist indicated that cohort studies with individual-level exposure-response data and the broader epidemiology literature both provide no evidence of increased mesothelioma risk due to chrysotile exposure. Further, this panelist noted that 33 of 41 mesothelioma cases previously identified as occurring among workers primarily exposed to chrysotile fibers (Stayner et al. 1996) were later reported as likely resulting from exposures to tremolite fibers found in the chrysotile mines (McDonald et al. 1997). This panelist noted that a recent finding of a small mesothelioma risk from chrysotile (Hodgson and Darnton 2000) results entirely on the assumption that the 33 mesothelioma cases mentioned above result entirely from chrysotile exposures. Based on these observations, this panelist indicated that the literature suggests that chrysotile exposures have limited, if any, role in causing mesothelioma. He nonetheless supported the relative potency attributed to chrysotile in the proposed protocol as a conservative measure in the overall risk assessment process.

Specific comments on the Connecticut friction products workers. Another panelist commented on an epidemiological study of a cohort of workers employed at a friction products plant in Connecticut. The panelist noted that the original study (McDonald et al. 1984) did not identify any deaths from mesothelioma, but review of the state cancer registry (Teta et al. 1983)

revealed that three Connecticut residents who died of mesothelioma were employed by the same friction products company. One of these employees had amphibole exposures during the time he worked for a textile plant that was under the same parent company that owned and operated the friction products plant. The other two cases, the panelist noted, were females who indeed worked at the friction products plant. A pathology review found that one of these cases was a woman with probable pleural mesothelioma and 5 years of exposure; the other case was a peritoneal mesothelioma in a woman who also had asbestosis, and worked as a clerk for 30 years. This panelist noted that it was questionable to attribute the latter two mesothelioma diagnoses to the chrysotile exposures at the friction products plant, though she added that this possibility cannot be definitively ruled out. This panelist encouraged that future review of this epidemiological study should be revised given this new information.

- # ***Comments on the proposed 500-fold difference in relative potency.*** The panelists had several comments on the finding in the proposed risk assessment methodology that amphibole fibers are 500 times more potent for mesothelioma than are chrysotile fibers. Several panelists noted that this finding is consistent with that of a recent re-analyses of 17 epidemiological studies (Hodgson and Darnton 2000). Though not disagreeing that amphibole fibers are clearly more potent, one panelist was concerned that the risk coefficients (K_M) were largely derived from data sets with inadequate exposure-response information for mesothelioma, and assumptions had to be made to determine critical inputs to the mesothelioma model (e.g., average exposure, duration of exposure).

Other panelists commented on specific sections in the proposed protocol. One panelist, for example, recommended that the authors check the accuracy of data presented in Table 6-16 and Table 6-29 of the report, which are not reported consistently. Another panelist suggested that the authors better explain why separate risk coefficients for amphiboles and chrysotile were calculated for some cohorts (e.g., Hughes et al. 1987) but not for others (e.g., Berry and Newhouse 1983), even though the exposure information available for the studies appears to be comparable. Finally, one panelist recommended that the authors of the proposed protocol consider questions recently raised (Rogers and Major 2002) about the quality of the exposure data originally reported for the Wittenoom cohort (De Klerk et al. 1989) when evaluating exposure-response relationships for mesothelioma.

3.2.2 Mesothelioma and Fiber Type: Inferences from Animal Toxicology and Mechanistic Studies

The panelists discussed the inferences provided by animal toxicology data and mechanistic data regarding relative mesothelioma potency of different asbestos fiber types. Overall, two panelists

commented that the human epidemiological data clearly establish that exposures to amphibole asbestos fibers pose a greater mesothelioma risk than do exposures to chrysotile fibers. They added that the animal toxicology data are generally supportive of this finding, but the animal data suffer from some limitations. Two panelists, for instance, noted that the utility of animal toxicology studies is limited by the fact that rodents are rather insensitive to mesothelioma. These panelists added that the animal toxicology studies involving intra-tracheal instillation or peritoneal injection are not directly relevant to the inhalation exposures that occur in humans. These limitations notwithstanding, the panelists raised the following points when discussing the animal toxicology and mechanistic studies:

One panelist referred to one of his earlier publications (Lippmann 1994) for further insights on the occurrence of mesothelioma in animal studies. At that time, this panelist noted, the animal inhalation studies found fewer than 10 cases of mesothelioma, and the number of cases appeared to be greatest among animals that were exposed to mixtures containing higher proportions of amphibole fibers. He found this consistent with the influence of fiber type observed in the human epidemiological data (see Section 3.2.1).

During this discussion, one panelist reviewed a publication (Suzuki and Yuen 2001) that was mentioned earlier in the workshop. The publication documents the amounts and types of asbestos fibers measured in samples of pleural plaques and tumor tissue collected for legal cases. These analyses reportedly found relatively large amounts of short, thin chrysotile fibers in the pleura, suggesting that these fibers should not be excluded from the group of fibers believed to induce mesothelioma. The panelist had several criticisms of the study. First, he indicated that the samples were analyzed using a non-standard technique, without any controls. Second, he questioned the major finding of fibers being detected in the pleura, because most of the samples analyzed were actually tumor tissue, in which he would not expect to find fibers. The panelist suspected that the chrysotile fibers reportedly found in the study likely result from specimen contamination—a bias that would have been more apparent had rigorous quality control procedures been followed. Finally, the panelist noted that a more rigorous study (Boutin et al. 1996) of

asbestos fibers in the parietal pleura found a mixture of fibers, including long amphibole fibers, among living patients with asbestos-related conditions. Based on these concerns, the panelist concluded that the publication of concern (Suzuki and Yuen 2001) is seriously flawed and its recommended should be excluded from EPA's analyses.

A specific issue raised regarding the analytical technique in the study (Suzuki and Yuen 2001) was that water was used during the digestion process. Noting that water may contain large amounts (>30,000 fibers/L) of small asbestos fibers, another panelist suspected that the fibers detected in the study might have resulted from contamination introduced during the digestion process. Because control samples were not analyzed, the panelist said the study offers no evidence that the fibers detected truly were in the original pleural plaques or tumor tissues. He added that studies of lung-retained asbestos fibers routinely detect primarily short, chrysotile fibers, and that the presence of the short fibers in the pleural tissue—even if the measurements from the study are valid—would not necessarily prove that short fibers cause mesothelioma.

3.2.3 Mesothelioma and Fiber Dimension: Inferences from the Epidemiology Literature

The panelists commented briefly on how the human epidemiological data characterize the role of fiber size on mesothelioma risk. Noting that exposure measurements in most every epidemiological study do not characterize fiber length distribution, one panelist indicated that these studies provide no direct evidence of how fiber length is related to mesothelioma. He added that the studies offer conflicting indirect evidence of the role of fiber length. Specifically, the higher mesothelioma risk coefficient among textile workers in South Carolina, when compared to that for the chrysotile miners and millers in Quebec, could be supportive of longer fibers being more potent, since exposures in South Carolina had a larger percentage of long fibers. However, a cohort of cement plant workers in New Orleans was found to have a higher mesothelioma risk coefficient than that of the South Carolina cohort, even though the South Carolina workers were exposed to higher percentages of long fibers. Finally, as indirect

evidence that carcinogenic potency increases with fiber length, this panelist noted that the mesothelioma risk model using the proposed exposure index, which is heavily weighted by long fibers, provided a considerably improved fit to the epidemiological data.

The panelists briefly revisited the inferences that can be drawn from studies of lung-retained fibers. One panelist again commented that results from a recent study (Suzuki and Yuen 2001) should be viewed with caution. He added that several other lung pathology studies (e.g., McDonald et al. 1989, Rogers et al. 1991, Rödelsperger et al. 1999) have been conducted using more rigorous methods, such as using appropriate controls for age, sex, and hospital. These studies all showed that risk of mesothelioma was considerably higher for individuals with larger amounts of long fibers retained in their lungs.

One panelist indicated that results from a study of lung-retained fibers (Timbrell et al. 1988) suggest fiber diameter plays a role in mesothelioma risk: the study observed no mesothelioma cases among a population highly exposed to anthophyllite fibers, which tend to be thicker fibers. Citing his earlier review of mesothelioma cases (Lippmann 1988), the panelist also noted that crocidolite fibers are both thinner than and more potent than amosite fibers, which further supports the hypothesis that carcinogenic potency for asbestos decreases with increasing fiber diameter.

3.2.4 Mesothelioma and Fiber Dimension: Inferences from Animal Toxicology and Mechanistic Studies

The panelists made few observations on findings from animal toxicology studies regarding mesothelioma and fiber length. One panelist indicated that findings from the animal toxicology studies generally support the overall finding that mesothelioma risks are greatest for long, thin fibers. However, another panelist noted that his earlier review of mesothelioma risks (Lippmann 1988) hypothesized that the critical fibers for mesothelioma induction are those with lengths between 5 and 10 μm . This panelist added that fibers of this dimension are more likely to translocate to the pleura than are longer fibers, but

he acknowledged that it is unclear whether fibers must first translocate to the pleura in order to cause mesothelioma.

Some panelists indicated that fiber durability likely plays a role in inducing mesothelioma, based on the fact that mesothelioma is more easily induced in animals using administration methods (e.g., peritoneal injection) that remove the importance of dissolution.

3.3 Exposure Estimates in the Epidemiology Literature

The panelists raised numerous issues when responding to the third charge question: “To what extent are the exposure estimates documented in the asbestos epidemiology literature reliable?” Recognizing that the exposure estimates from the epidemiology studies are critical inputs to the exposure-response assessment, the panelists expressed concern about the exposure data: few studies provide detailed information on fiber size distribution; many studies report exposures using outdated sampling and analytical methodologies (e.g., midget impinger); individual-level data are not available for most studies; and many studies do not report detailed information on parameters (e.g., exposure levels, exposure duration) needed to evaluate exposure-response relationships, particularly for mesothelioma. Their specific concerns on these and other matters follow:

- # ***Concerns regarding exposure estimates in specific studies.*** Some panelists expressed concern about the assumptions made to interpret the exposure data originally reported in the epidemiology studies. One panelist reviewed specific examples of these concerns:
- ▶ The original study of workers at a Connecticut friction products plant (McDonald et al. 1984) reports exposures measured by midget impingers (in units of mmpcf), with no information on how to convert this to PCM measurements, and the original publication includes limited data on exposure duration.
 - ▶ The original study of workers at a New Jersey insulation factory (Seidman et al. 1986) did not report any exposure measurements from the factory studied, and data collected

from another plant with similar operations were used to characterize exposure-response for this cohort.

- ▶ The original study of workers at a Texas insulation factory (Levin et al. 1998) reported a range of exposure levels (15–91 fibers/mL), and the authors of the proposed protocol assigned an average exposure level (45 fibers/mL) to the entire cohort.
- ▶ The original study of U.S. insulation applicators (Selikoff and Seidman 1991) has no information on exposure. The proposed protocol assumes that all workers were exposed to 15 fibers/mL for 25 years, based on a separate review of exposures among insulation workers (Nicholson 1976).
- ▶ The original study of retirees from the U.S. Asbestos Products Company (Enterline et al. 1986) reported exposures based on midget impinger sampling, with no information on how to convert these exposures to PCM measurements.
- ▶ According to a recent letter to the editor (Rogers and Major 2002), the original study of the Wittenoom cohort (De Klerk et al. 1989) might have overestimated exposures, possibly by as much as a factor of 10.

The previous comments led to a discussion on whether certain studies should be excluded from the meta-analysis used in the proposed protocol (see next bulleted item). Prior to this discussion, one panelist expressed concern about being overly critical of the exposure estimates used for many of the studies listed above; he emphasized that all exposure estimates appear to be based on a critical review of the literature, and no estimates are completely arbitrary, as some of the panelists' comments implied.

Comments on using study inclusion criteria for the meta analysis. Given the concerns about the quality of exposure data reported in some epidemiology studies, the panelists debated whether future revisions of the proposed protocol should exclude certain studies from the exposure-response analysis. The panelists were divided on this matter.

On the one hand, several panelists recommended that the authors develop and apply study inclusion criteria in the exposure-response evaluation, as is commonly done when conducting a meta-analysis. One panelist, for instance, recommended assessing exposure-response relationships for only those studies found to have adequate exposure data, and then using a sensitivity analysis to examine the effect of excluding studies with inadequate exposure data. These panelists clarified that they are not advocating disregarding the majority of studies; rather, they are suggesting simply that the authors of the proposed protocol use study inclusion criteria and sensitivity analyses to ensure that the conclusions are based on the best available exposure data.

On the other hand, several panelists supported the current approach of using as many studies as possible and accounting for the quality of the exposure measurements in the uncertainty factors. One panelist, for example, commended the authors for being as inclusive as possible when reviewing the studies; he supported the approach of recognizing the limitations of the available exposure data and accounting for these limitations in the uncertainty factors that were ultimately used to weight the studies in the meta-analysis. This panelist acknowledged that the exposure estimates in some of the epidemiological studies might be rough estimates, but he emphasized that the estimates are not worthless and should not be discarded. Other panelists concurred with these comments, and did not support applying overly restrictive study inclusion criteria.

Comments on the uncertainty factors assigned to each study. The panelists made several comments on the uncertainty factors that the authors assigned to each study. Dr. Berman first explained the four uncertainty factors: the first factor (F1) characterizes the confidence in exposure estimates; the second factor (F2) represents the confidence in the conversion to PCM measurements from other exposure metrics (typically midge impinger analyses); the third factor (F3) characterizes the confidence the authors had on worker history data; and the fourth factor (F4) was a non-exposure related factor to account for other uncertainties (e.g., lack of information on confounders, incomplete or inaccurate mortality ascertainment). Dr. Berman described generally how the individual uncertainty factors were assigned and noted that each factor could range from 1 to 5.

The panelists' comments primarily focused on the transparency of how uncertainty factors were presented and incorporated into the meta-analysis. Multiple panelists, for instance, recommended that future revisions to the proposed protocol include a table that lists the uncertainty factors assigned to each study. Further, one panelist suggested that the revised protocol describe the assumptions inherent in the uncertainty factor weighting approach, such as explaining why some factors are assigned values over a broader range than others (e.g., why F1 values span a broader range than F4 values) and describing why the individual uncertainty factors have equal weights in generating the composite uncertainty factor. Another panelist agreed, and added that the revised protocol should more explicitly describe how the uncertainty factors were combined into the composite factor and how this composite factors affects the weighting of studies in the meta-analysis. Expanding on this point, another panelist suggested that the final document more clearly explain that the final estimates of cancer risk coefficients (K_L^* and K_M^*) are actually weighted averages of the epidemiological studies, with the weights assigned to each study being a function of that study's uncertainty. This panelist also recommended that the revised document clearly state how, if at all, the fraction of amphibole fibers and the fraction of fibers longer than 10 μm are reflected in the uncertainty factors.

Some panelists debated the utility of alternate approaches that could be used to assign uncertainty factors. Two panelists noted that the approach used to assigning uncertainty factors is somewhat subjective, because different groups of analysts would likely assign different

uncertainty factors. To avoid the appearance of arbitrariness, these panelists suggested using alternate meta-analysis approaches that do not require using uncertainty factors. They noted, for example, that the authors could use a random effects model in which residual inter-study variation is estimated. Another suggestion was to conduct sensitivity analyses examining the effects of including or excluding studies, depending on the uncertainty factors assigned to them.

Another panelist disagreed with these comments and supported the analyses in the proposed protocol; this panelist indicated that the authors had no choice but to make judgments based on the information documented in the epidemiology literature. He suggested that EPA consider convening a separate expert panel to assign uncertainty factors, if panelists do not support those selected by Drs. Berman and Crump.

Assumptions made to convert exposure estimates from midget impinger sampling.

Several panelists noted that the original publications for many epidemiology studies document exposure estimates based only on midget impinger sampling and do not include any information on how to convert these exposures to levels that would be measured by more modern methods (e.g., PCM, TEM). The panelists noted that the conversion factor (from mmpcf to fibers/mL) can vary considerably from one occupational setting to the next.

Interpretations of the study of South Carolina textile workers. The panelists had different opinions on interpretations of the study of South Carolina textile workers (Dement et al. 1994). One panelist, for instance, found this particular study to be an outlier among the other epidemiological studies, and he recommended that the authors exclude this study from the exposure-response analysis until the causes for the increased relative risks observed for this cohort are better understood. Another panelist suggested that the proposed protocol should classify the South Carolina cohort as being exposed to mixed asbestos fibers, rather than being exposed to chrysotile fibers. He indicated that some workers in the cohort were exposed to amosite and crocidolite, in addition to being exposed to chrysotile.¹

Other panelists, however, did not think the South Carolina study should be excluded from EPA's analysis. One panelist was troubled about criticisms of the exposure estimates for this cohort, given that this is one of few studies in which co-located samples were collected and analyzed using different methods, thus providing site-specific data for converting midget impinger

¹ After reviewing a draft of this report, one panelist indicated that it is important to note that exposure data for the South Carolina cohort are available from more than just one reference (Dement et al. 1994). He suggested that EPA use data from studies conducted by McDonald in the 1980s of a parallel cohort in the same plant. However, he cautioned EPA against treating multiple studies of the same relatively small group of workers as separate studies, considering the large overlap of workers studied by the two groups of investigators. This panelist encouraged EPA to consider other data sources for this cohort, given that a recent re-analysis of epidemiological studies (Hodgson and Darnton 2000) severely criticized the data source EPA uses (Dement et al. 1994), to the point of those data being dropped from the recent re-analysis altogether.

sampling results to PCM measurements. Another panelist challenged suggestions that the South Carolina study is an outlier; he indicated that the South Carolina study is one of the more rigorous epidemiology studies available for asbestos exposures, and he found no valid scientific reasons for discarding it. During this discussion, one panelist point out in response that the South Carolina study is indeed an outlier among the textile cohorts, with a slope which is higher than either of the two textile cohorts; this panelist did acknowledge that the lung cancer risk among the textile cohorts is greater than that among the mining cohorts. This panelist added that scientists need a better explanation for why the lung cancer risk among the South Carolina cohort is greater than that of other cohorts before the South Carolina study can achieve credibility, especially considering that exposures in South Carolina were supposedly to “pure” chrysotile.

4. COMMENTS ON TOPIC AREA 2: THE PROPOSED EXPOSURE INDEX

This section summarizes the panelists' responses to the charge questions pertaining to the proposed exposure index. Section 4.1, 4.2, and 4.3 document the panelists' responses to charge questions 4, 5, and 6, respectively.

4.1 Responses to Charge Question 4

Charge question 4 asks: "The proposed exposure index does not include contributions from fibers shorter than 5 μm . Please comment on whether the epidemiology and toxicology literature support the conclusion that asbestos fibers shorter than 5 μm present little or no carcinogenic risk." The panelists discussed this matter earlier in the workshop (see Sections 3.1.3 and 3.1.4 for these comments), and provided additional insights on the matter. Overall, the panelists agreed that carcinogenic potency increases with fiber length, particularly for lung cancer. Most panelists supported assigning no potency to fibrous structures smaller than 5 μm . Some panelists agreed that the short fibrous structures are clearly less potent than long fibers, but they had reservations about assigning zero potency to the structures smaller than 5 μm ; these panelists acknowledged that the toxicity of the short fibrous structures might be adequately addressed by EPA's air quality standards for particulate matter. Specific comments on this charge question follow:

- # ***Reference to ATSDR's expert panel workshop on the role of fiber length.*** Two panelists noted that ATSDR convened an expert panel in October 2002 to discuss the role of fiber length on toxicity, and much of that discussion specifically addressed fibrous structures smaller than 5 μm . A main conclusion of that panel was that there is "a strong weight of evidence that asbestos and synthetic vitreous fibers shorter than 5 μm are unlikely to cause cancer in humans" (ERG 2003). The panelists encouraged EPA to review the summary report prepared for that workshop, which was officially released on March 17, 2003, and is available on-line at: www.atsdr.cdc.gov/HAC/asbestospanel.
- # ***Evidence from epidemiological studies.*** One panelist indicated that the epidemiological studies do not provide direct evidence of the role of fibrous structures shorter than 5 μm .

However, the panelist indicated that a growing body of evidence suggests that the cohorts predominantly exposed to shorter fibers (e.g., friction brake workers, gold miners, taconite miners) do not have statistically significant increased cancer risks. This panelist added that the mechanistic studies provide the strongest evidence for assigning no potency to fibrous structures (see next bulleted item). Another panelist agreed with these statements, and added that his interpretation of data compiled by the National Cancer Institute provide additional indirect evidence of short fibrous structures presenting little or no carcinogenic risk (see page 102 of the premeeting comments in Appendix B).

The panelists briefly revisited the findings from a recent publication (Suzuki and Yuen 2001) that reported finding relatively large amounts of short, thin chrysotile fibers in malignant mesothelioma tissue. Several panelists encouraged that these findings not be considered in the risk assessment methodology for reasons cited earlier in the workshop (see Section 3.2.2).

- # ***Evidence from mechanistic studies.*** The panelists offered different interpretations of mechanistic studies. One panelist indicated that mechanistic studies have shown that shorter fibers are cleared more readily than long fibers from the alveolar region of the lung by phagocytosis, and therefore provide supporting evidence that short fibers play little or no role in carcinogenic risk. This panelist acknowledged that extremely high doses of particular matter and other non-fibrous structures can generate biological responses (e.g., inflammation), but he doubted that such “overload” conditions would be relevant to the environmental exposures that the proposed protocol will be used to evaluate.

Another panelist agreed that long fibers are clearly more potent than short fibrous structures, but he questioned the conclusion that short fibrous structures have no impact on carcinogenic risk. This panelist noted that mechanistic studies have demonstrated that short fibrous structures and spherical particles, like silica, can elicit the same toxic responses (e.g., generate reactive species, stimulate proliferative factors) identified for asbestos fibers. This panelist added, referring to his premeeting comments, that exposure to short fibers could cause inflammation and generation of oxidative species that might increase the response to long fibers (see Bellman et al. 2001). Overall, this panelist acknowledged that long fibers are more persistent than short fibers in the lung and should be weighted more heavily in the exposure index, but he was hesitant to assign the short fibrous structures zero potency.

- # ***Implications on sampling and analytical methods.*** One panelist commented on the practical implications, from a sampling perspective, of any changes to the exposure index. This panelist indicated that measuring all fibers (including structures shorter than 5 μm) in environmental samples would not only be expensive, but also would compromise the sensitivity of measuring the longer fibers that are most predictive of cancer risk. This panelist acknowledged that human exposure is predominantly to fibrous structures less than 5 μm , but he noted that the amounts of short fibrous structures retained by the lung tend to be very strongly

correlated with the amounts of long fibers retained by the lung. Due to this correlation, this panelist noted that measuring long fibers with sufficient accuracy would allow one to estimate amounts of short fibrous structures in a sample. This panelist added, however, that he sees no benefit of characterizing exposures to fibrous structures smaller than 5 μm , given the conclusion that such fibers do not cause cancer (ERG 2003).

4.2 Responses to Charge Question 5

Charge question 5 asks: “The proposed exposure index is weighed heavily by fibers longer than 10 μm . Specifically, Equation 7.13 suggests that the carcinogenic potency of fibers longer than 10 μm is more than 300 times greater than that of fibers with lengths between 5 and 10 μm . How consistent is this difference in carcinogenic potency with the epidemiology and toxicology literature?” The panelists’ responses to this question follow:

- # ***Consistency with epidemiological literature.*** The panelists noted that the original epidemiology studies did not collect exposure information that provides direct evidence of the relative potency assigned to the two different fiber length categories: fibers longer than 10 μm , and fibers with lengths between 5 and 10 μm . During this discussion, one panelist recommended that EPA consider the results of a case-control study (Rogers et al. 1991) that suggests that mesothelioma risks are greater for individuals with larger amounts of the shorter fibers (i.e., between 5 and 10 μm) retained in their lungs. Another panelist was not convinced of the findings from this study, due to possible biases from selection of controls not matched for hospital of origin. This panelist encouraged EPA to refer to more rigorous lung-retained fiber studies (e.g., McDonald et al. 1989, Rödelsperger et al. 1999) that have found that the majority of cancer risk for mesothelioma is attributed to exposures to longer fibers, even when measurements of short fibers are taken into account.
- # ***Questions about the fiber length-dependence used for mesothelioma.*** Some panelists were not convinced that the relative potencies assigned to different fiber lengths were appropriate for mesothelioma. One panelist, for instance, noted that his previous review of the literature (Lippmann 1994) suggests that cancer risk for mesothelioma is most closely associated with exposure to fibers between 5 and 10 μm long. He indicated that this assessment is consistent with other human lung evaluations (e.g., Timbrell et al. 1988), which have reported that fibers retained by the lung tend to be longer than fibers that translocate to the pleura. This panelist added that the epidemiology literature clearly suggests that lung cancer and

mesothelioma have different risk factors, as the relative amounts of lung cancer and mesothelioma cases vary considerably from one cohort to the next. Based on these concerns, this panelist suggested that EPA consider developing separate fiber length weighting schemes for lung cancer and mesothelioma.

Another panelist indicated that the epidemiology studies provide indirect evidence that carcinogenic potency appears to increase with fiber length. Specifically, he noted that the studies consistently show that mesothelioma has a very long latency period—a trend that suggests that the most durable fibers (i.e., the longer fibers) are the most potent. The panelist added that the analyses in the proposed protocol provide further indirect evidence of mesothelioma risks increasing with fiber length: when the exposure index was used in the mesothelioma model, the proposed risk assessment methodology generated an improved fit to the epidemiological data.

During this discussion, a panelist cautioned about inferring that only those fibers that reach the pleura are capable of causing mesothelioma, because researchers have not determined the exact mechanisms by which mesothelioma is induced. Further, he cautioned about inferring too much from a single study (Timbrell et al. 1988), given that many additional studies are available on lung-retained fibers.

Questions about the relevance of animal toxicology data. Some panelists expressed concern about basing the proposed weighting factors for different fiber lengths on observations from animal data. First, one panelist noted that the weighting factors were derived strictly based on lung cancers observed in laboratory animals, and he questioned whether one can assume that the weighting factors can be defensibly applied to mesothelioma. Second, other panelists noted that extrapolating the weighting factors from rodents to humans also involves uncertainty, due to inter-species differences in respiratory anatomy, macrophage sizes, and sites of lung cancers.

Suggested follow-up analyses. Given the concerns about basing the proposed exposure index entirely on data from animal toxicology studies, two panelists recommended that EPA attempt to optimize the weighting factors applied to different fiber length categories using the available human epidemiological data. One panelist suggested that this optimization could be performed using the data compiled in Table 6-15 in the proposed protocol, which presents estimates of the fiber length distribution for different occupational cohorts. A panelist also suggested that EPA consider deriving separate weighting factors for lung cancer and mesothelioma, rather than assuming the same fiber length dependence for both outcomes.

4.3 Responses to Charge Question 6

Charge question 6 asks: “Please explain whether the proposed exposure index will allow meaningful comparisons between current environmental exposures to asbestos and historical exposures to asbestos that occurred in the work place.” The panelists discussed several topics when addressing the question, because some panelists had different impressions of what the question was asking. Some panelists viewed the question as asking about the validity of low-dose linear extrapolations (see Section 3.1.5 for more information on this topic), and others viewed the question as asking about whether the proposed methodology is an improvement over EPA’s current risk assessment model. A summary of the panelists’ specific responses follows:

Is the proposed exposure index an improvement to asbestos risk assessment? When answering this charge question, multiple panelists focused on whether the proposed exposure index is an improvement over EPA’s 1986 asbestos risk models. These panelists agreed that the proposed approach is more consistent with the overall literature on health risks from asbestos, which show that cancer risks vary with fiber type and fiber dimension. Two panelists were hesitant to call the proposed approach an improvement for evaluating mesothelioma risks, because the fiber length weighting factors are based entirely on lung cancer data in animals. These panelists were particularly concerned that the proposed methodology might assign lower risks for mesothelioma in certain circumstances, because the fiber-length dependence in the methodology is not based on any toxicological or epidemiological studies of mesothelioma.

Does the proposed risk assessment model support extrapolation from occupational exposures to environmental exposures? Some panelists commented on the applicability of the proposed risk assessment model to exposure doses below the ranges considered in the occupational studies. Referring to observer comments provided earlier in the workshop, two panelists indicated that some environmental exposures in areas with naturally-occurring asbestos do not appear to be considerably lower than those experienced by occupational cohorts. Another panelist agreed, and cautioned about distinguishing environmental exposures from occupational exposures; he instead encouraged EPA and the panelists to focus on the exposure magnitude, regardless of whether it was experienced in an occupational or environmental setting.

One panelist recommended that EPA investigate how cancer risks for lung cancer and mesothelioma vary between EPA’s 1986 model and the proposed risk assessment methodology: for different distributions of fiber types and dimensions, does the proposed methodology predict higher or lower risks than the 1986 model? Dr. Berman indicated that the proposed methodology, when compared to EPA’s 1986 model, generally predicts substantially higher risks for environments with longer, thinner fibers and environments with larger amounts of

amphibole fibers and predicts somewhat lower risks for environments with shorter, thicker fibers and environments that contain only chrysotile fibers. One panelist recommended that future revisions to the proposed protocol include sample calculations, perhaps in an appendix, for several hypothetical environments to demonstrate how estimated cancer risks compare between the new methodology and the 1986 model.

5. COMMENTS ON TOPIC AREA 3: GENERAL QUESTIONS

This section summarizes the panelists' responses to charge questions 7–10 and 12. Responses to charge question 11 are included in Section 6, because this charge question sought the panelists' overall impressions of the proposed risk assessment methodology, rather than focusing on any one specific issue.

5.1 Responses to Charge Question 7

This charge question asks: “The proposed risk assessment approach assigns carcinogenic potency to individual fibers and to cleavage fragments (or ‘bundles that are components of more complex structures’). Please comment on whether cleavage fragments of asbestos are as toxicologically significant as fibers of the same size range.” The panelists raised the following points when responding:

- # ***Terminology used in the charge question.*** One panelist took strong exception to the wording in this question (see pages 30–33 in Appendix B) and strongly recommended that the panelists use correct terminology during their discussions. This panelist noted, for instance, that cleavage fragments are not equivalent to bundles, nor do cleavage fragments meet the regulatory definition of asbestos, as the charge question implies. He clarified that he defines cleavage fragments as non-asbestiform amphiboles that are derived from massive amphibole structures. This panelist was concerned that none of the panelists at the workshop has the mineralogical expertise needed to address issues pertaining to cleavage fragments. Another panelist echoed these concerns and agreed that this charge question raises complex issues.

- # ***Significance of cleavage fragments with respect to human health effects.*** The previous concerns notwithstanding, several panelists commented on the role of cleavage fragments in the proposed risk assessment methodology. One panelist, for example, indicated that there is no reason to believe that cleavage fragments would behave any differently in the human lung than asbestiform fibers of the same dimensions and durability; he added that this conclusion was also reached by the American Thoracic Society Committee in 1990 (Weill et al. 1990). This panelist acknowledged, however, that expert mineralogists have differing opinions on the role of cleavage fragments. Several other panelists agreed that it is reasonable to assume that cleavage fragments and asbestos fibers of the same dimension and durability would elicit similar toxic responses.

- # ***Review of selected epidemiological and toxicological studies.*** The panelists briefly discussed what information has been published on the toxicity of cleavage fragments. One panelist indicated that Appendix B in the proposed protocol (see pages B-3 through B-10) interprets results from an animal study (Davis et al. 1991) that evaluated exposures to six tremolite samples, including some that were primarily cleavage fragments. This panelist noted that the study provides evidence that cleavage fragments can cause mesothelioma in animals.

Another panelist, however, cautioned against inferring too much from this animal study for several reasons: the study was not peer reviewed; the fiber measurements in the study reportedly suffered from poor reproducibility; and the mesotheliomas observed in the study might have reflected use of intra-peritoneal injection model as the dose administration method. This panelist recommended that EPA conduct a more detailed review on the few studies that have examined the toxicity of cleavage fragments, possibly considering epidemiological studies of taconite miners from Minnesota (Higgins et al. 1983) and cummingtonite-grunerite miners from South Dakota (McDonald et al. 1978); he noted that a pending publication presents updated risks among the taconite miners.

- # ***Practical implications of measuring cleavage fragments in environmental samples.*** One panelist added, and another agreed, that measuring cleavage fragments in environmental samples presents some challenges, because microscopists cannot consistently distinguish cleavage fragments from asbestiform fibers, even when using TEM.

5.2 Responses to Charge Question 8

Charge question 8 asks: “Please comment on whether the proposed cancer assessment approach is relevant to all amphibole fibers or only to the five types of amphibole fibers (actinolite, amosite, anthophyllite, crocidolite, tremolite) designated in federal regulations.” The panelists made the following general comments in response:

- # ***Review of evidence from toxicological and epidemiological studies.*** The panelists identified few studies that address the toxicity of amphibole fibers other than actinolite, amosite, anthophyllite, crocidolite, and tremolite. One panelist indicated that animal toxicology studies have demonstrated that synthetic vitreous fibers with differing chemistry, but having similar durability and dimensions, generally exhibit similar potency for fibrosis, lung cancer, and mesothelioma. Another panelist added that lung cancer and mesothelioma exposure-response

relationships for a cohort of vermiculite miners from Libby, Montana, have been published for both asbestiform richterite and winchite.

- # ***Appropriateness of applying the model to non-asbestiform amphiboles.*** Several panelists agreed that the proposed risk assessment methodology is relevant to amphibole fibers other than those listed in the federal regulations. The panelists noted that, in the absence of more detailed information on the matter, it is prudent to assume that fibers of similar dimension and durability will exhibit similar toxic effects. Two panelists expressed some hesitation on applying the proposed model to the non-asbestiform amphiboles: one panelist asked how confidently one can apply the cancer risk coefficients to amphibole fibers that have not been studied, and another panelist indicated he was not convinced that the model should be applied to the other amphiboles, let alone for the amphiboles that are listed in the federal regulations.

Given the amount of naturally occurring amphiboles in the Earth's crust, one panelist suggested that the proposed protocol clearly state that the non-asbestiform amphiboles being evaluated are only those with the same dimensional characteristics and biodurability as the corresponding asbestiform amphiboles.

5.3 Responses to Charge Question 9

Charge question 9 asks: "The review document recommends that asbestos samples be analyzed by transmission electron microscopy (TEM) and count only those fibers (or bundles) longer than 5 μm . Such counting practices will provide no information on the amount of asbestos fibers shorter than 5 μm . To what extent would data on shorter fibers in samples be useful for future evaluations (e.g., validation of the cancer risk assessment methodology, assessment of non-cancer endpoints)?"

The panelists expressed varying opinions on this matter: some panelists saw no benefit of measuring fibrous structures shorter than 5 μm , based on responses to earlier charge questions (see Sections 3.1.3, 3.1.4, and 4.1); other panelists indicated that there is some utility to collecting information on shorter fibrous structures, particularly if the incremental analytical costs are not prohibitively expensive and if counting short fibers does not compromise accurate counts of longer fibers. The panelists raised the following specific issues when discussing measurement methods:

- # ***Support for using TEM in future sampling efforts.*** The panelists unanimously supported the recommendation in the proposed protocol of using TEM, rather than PCM or some other method, to characterize exposures in future risk assessments. The panelists also emphasized that future measurement methodologies must focus on generating accurate counts of the most biologically active fibers, or fibers longer than 5 μm .
- # ***Practical implications of counting fibers shorter than 5 μm .*** One panelist indicated that analyzing samples for fibrous structures shorter than 5 μm would compromise analysts' ability to accurately count the amounts of longer fibers that are of greater biological concern. Some panelists and an observer further discussed the costs associated with counting fibers in multiple length categories, including shorter than 5 μm . The panelists did not cite firm cost figures for these analyses. However, noting that environmental samples typically contain more than 90% short fibrous structures, one panelist suspected that counting the shorter structures would considerably increase the time a microscopist needs to analyze samples, and therefore also would considerably increase the cost of the analysis. A panelist indicated that the costs and benefits of counting fibers shorter than 5 μm might be more appropriately debated between microscopists and risk assessors, with inputs from industrial hygienists and mineralogists.
- # ***Relevance of fibers shorter than 5 μm for non-cancer endpoints.*** One panelist noted that exposures to fibrous structures shorter than 5 μm can contribute to asbestosis in occupationally exposed individuals (Lippmann 1988), but he doubted that the exposure levels found to be associated with asbestosis would be experienced in non-occupational settings. Another panelist added that the role of shorter fibrous structures for other non-cancer endpoints is not known, such as the pleural abnormalities and active pleural fibrosis observed in Libby, Montana. No panelists were aware of any authoritative statements made on the role that short fibers play, if any, on these other non-cancer endpoints. During this discussion, one panelist indicated that the toxicity of fibrous structures shorter than 5 μm might be adequately addressed by EPA's particulate matter standards.

5.4 Responses to Charge Question 10

Charge question 10 asks: "The proposed risk assessment methodology suggests that exposure estimates should be based only on fibers longer than 5 μm and thinner than 0.5 μm . Is this cut-off for fiber diameter appropriate?" Before the panelists responded to the question, Dr. Berman first clarified that the exposure index optimized from the animal studies (see Equation 7.12 in the proposed protocol)

assigns a far greater carcinogenic potency to fibers longer than 40 μm , with diameters less than 0.4 μm ; he noted that the proposed diameter cut-off (0.5 μm) was based on an *ad hoc* adjustment.

The panelists agreed that the proposed cut-off for fiber diameter (0.5 μm) would likely include most fibers of health concern; however, they also unanimously agreed that the exposure index should not exclude thicker fibers that are known to be respirable in humans. The main argument given for increasing the cut-off is that fibers with diameters as large as 1.5 μm (or with aerodynamic diameters as large as 4.5 μm) can penetrate to small lung airways in humans. Other panelists provided additional specific comments, generally supporting inclusion of thicker fibers in the proposed exposure index. One panelist, for example, advised against basing the fiber diameter cut-off strictly on observations from rat inhalation studies, due to inter-species differences in respirability. Further, noting that the proposed cut-off for fiber diameter would likely exclude some amosite fibers and a considerable portion of tremolite fibers with known carcinogenic potency, another panelist encouraged that the proposed exposure index include contributions from thicker fibers.

The panelists noted that consideration of fibers thicker than 0.5 μm was viewed as being most important for the lung cancer risk assessment model, as risks for mesothelioma appear to be more closely linked to exposures to long, thin fibers (see Section 3.2.3). Further, some panelists suspected that increasing the fiber diameter cut-off in the exposure index should be accompanied by changes to the exposure-response coefficients in the risk assessment models, but the panelists did not unanimously agree on this issue.

5.5. Responses to Charge Question 12

Charge question 12 asks: “Section 8.2 of the review document presents three options for assessing cancer risks from asbestos exposure. Please comment on the technical merit of the proposed risk assessment options.” The panelists briefly reviewed the strengths and weaknesses of the three options presented in the proposed protocol for assessing asbestos-related cancer risks. The panelists agreed that the first option—direct use of EPA’s lung cancer and mesothelioma risk assessment models—allows for the greatest flexibility in evaluating site-specific exposure scenarios, particularly those with time-varying exposures. Dr. Crump indicated that he envisioned this option being coded into a computer program, into which users enter their site-specific exposure information. Most panelists endorsed developing such a program. The panelists did not reject use of the second and third options, provided that EPA ensures that all three options generate equivalent risk estimates for the same exposure scenario.

The one issue discussed in greater detail was how sensitive predictions using the first option are to the mortality rates used in the evaluation. Noting that mortality rates as functions of age and sex differ from one location to the next, this panelist encouraged EPA to consider carefully whether nationwide mortality estimates would be programmed into the risk assessment model or whether risk assessors would have the option of entering site-specific mortality rates. The panelist also suggested that the authors of the risk assessment conduct sensitivity analyses to quantify how strongly the mortality data affect cancer risk estimates. These comments also raised questions about the fact that two populations with different underlying mortality rates could have different cancer risks, even though their asbestos exposure levels are equivalent.

6. COMMENTS ON TOPIC AREA 4: CONCLUSIONS AND RECOMMENDATIONS

This section reviews the panelists' individual conclusions and recommendations regarding the proposed protocol (Section 6.1), as well as how the panelists developed their overall conclusions and recommendations that appear in the executive summary of this report (Section 6.2).

6.1 Responses to Charge Question 11

Charge question 11 asks: "Discuss whether the proposed cancer assessment approach, as a whole, is a reasonable evaluation of the available health effects data. What aspects of the proposed cancer assessment approach, if any, are inconsistent with the epidemiology or toxicology literature for asbestos?" The panelists offered individual summary statements, which were not discussed or debated among the panel. Following is a summary of the panelists' individual summary statements in the order they were given:

- # ***Dr. Lippmann's summary statement.*** Dr. Lippmann commended Drs. Berman and Crump on developing the proposed risk assessment protocol and supported use of a model that accounts for the factors (e.g., fiber type and dimension) that are most predictive of cancer risk. Dr. Lippmann supported the authors' attempt to make full use of the existing data and to interpret the results from the epidemiological studies. He strongly recommended that EPA make every effort to obtain individual-level data from additional epidemiological studies. Dr. Lippmann suggested that a follow-up workshop with experts in exposure assessment could help EPA evaluate the uncertainties in exposure measurements from historic occupational data sets. Dr. Lippmann supported an observer's suggestion to conduct an animal inhalation study using tremolite cleavage fragments to help resolve the issue of these fragments' carcinogenic potency. Overall, he encouraged that future work on the proposed protocol continue, through use of additional expert panels, to make more informed usage of the human exposure data.

- # ***Dr. Teta's summary statement.*** Dr. Teta indicated that the proposed protocol is an impressive integration of the animal toxicology data and the human epidemiology data. She commended the authors for developing a scientific methodology that successfully reduces the variability in results across the epidemiological studies, suggesting that the studies might be more consistent than were previously thought. Dr. Teta recommended improvements to the meta-analysis of

epidemiological studies, such as establishing and applying criteria for use of human data in characterizing exposure-response relationships. Overall, Dr. Teta found no inconsistencies between the proposed protocol and the larger body of epidemiology literature, including studies of cohorts (e.g., gas mask workers, railroad workers, friction brake workers) that do not have well-defined exposure information. Though not disagreeing with the utility of other panelists' recommendations, such as re-analyzing data from additional epidemiological studies and convening additional expert panels, Dr. Teta encouraged EPA to move forward expeditiously with completing the proposed protocol and discouraged implementing additional steps that might delay the overall project.

Dr. Hoel's summary statement. Dr. Hoel encouraged the use of more sophisticated modeling that incorporates data on exposure-response (including non-linear models), duration of exposure, cessation of exposure, and uncertainty in exposure. Dr. Hoel also strongly recommended that EPA attempt to obtain individual-level data from additional epidemiology studies, or at least obtain partial data sets. He encouraged Drs. Berman and Crump to use more sophisticated uncertainty analysis techniques, such as generating prior and posterior distributions of uncertainty. To ensure that the lung cancer model is not confounded by cigarette smoking, Dr. Hoel recommended that Drs. Berman and Crump more closely evaluate all available data on the interactions between asbestos exposure and cigarette smoking.

Dr. Steenland's summary statement. Dr. Steenland indicated that the proposed protocol is a step forward in asbestos risk assessment; however, he had several recommendations for improving the analysis of epidemiological studies. For instance, Dr. Steenland suggested that the authors conduct meta-regression analyses using the original exposure-response coefficients, in which predictor variables include fiber size, fiber type, the estimated percentage of amphiboles, percentage of fiber greater than 10 μm , and categorical grouping of studies according to quality. He indicated that these factors can be examined using both fixed effects and random effects models. Dr. Steenland recommended that the proposed protocol explicitly state and defend the basis for choosing the 10 μm cut-off for fiber length in the exposure index. He suggested that EPA should consider using Bayesian techniques or other methods to determine which relative potencies assigned to different fiber length categories optimize the model's fit to the epidemiological data.

Focusing on specific topics, Dr. Steenland indicated that he disagrees with the approach of assigning amphibole fibers five times greater lung cancer potency than chrysotile fibers, especially considering that the statistical analysis in the proposed protocol could not reject the hypothesis that amphibole fibers and chrysotile fibers are equally potent. Further, he advocated suggestions of exploring the adequacy of other exposure-response models (e.g., non-linear models). Finally, Dr. Steenland suspected that cigarette smoking likely will not be a confounding factor in exposure-response analyses for two reasons. First, he noted that differences in smoking practices between working populations and general populations typically do not cause

substantial differences in standardized mortality ratios. Second, he indicated that it is highly unlikely that prevalence of smoking varies with workers' exposure levels. Dr. Steenland encouraged that EPA refer to a recent publication (Liddell and Armstrong 2002) for similar insights on interactions between asbestos exposure and cigarette smoking.

- # ***Dr. Crapo's summary statement.*** Dr. Crapo complimented Drs. Berman and Crump on preparing the cancer risk assessment methodology, and he supported the general approach of expressing cancer risk as a function of asbestos fiber type and fiber dimension. Dr. Crapo indicated that the proposed protocol reaches several defensible conclusions, such as assigning greater mesothelioma potency to amphibole fibers and to longer fibers while assigning no risk to fibers less than 5 μm in length. However, he was concerned about some specific issues that are not yet adequately resolved. For instance, Dr. Crapo felt additional data are needed to rigorously define how mesothelioma potency varies with fiber length (i.e., fibers longer than 10 μm being 300 times more potent than fibers with lengths between 5 and 10 μm). Dr. Crapo recommended that EPA, when revising the proposed protocol, explore more sophisticated modeling techniques, including non-linear exposure-response models and consideration threshold effects. He supported more detailed analyses of interactions between asbestos exposure and cigarette smoking, again through the use of non-linear models.

- # ***Dr. Sherman's summary statement.*** Dr. Sherman first indicated that she concurred with several recommendations made by Drs. Hoel and Steenland. She focused her summary statements on the proposed exposure index, recommending that Drs. Berman and Crump use the epidemiology data to further investigate other formulations of an exposure index. Dr. Sherman recommended, for example, examining the goodness of fit of other formulations of the exposure index (e.g., assigning zero potency to all fibers shorter than 10 μm). Further, she recommended that the authors attempt to optimize the potency weighting factors in the exposure index to the epidemiological data. Finally, given that panelists expressed concern regarding how potency varies with fiber length for mesothelioma, Dr. Sherman suggested that Drs. Berman and Crump consider developing two different exposure indexes—one optimized for lung cancer, and the other for mesothelioma. Dr. Sherman added that she generally supported the lung cancer and mesothelioma exposure-response models, and questioned whether using more complicated models would necessarily lead to a better understanding of the data.

- # ***Dr. Castranova's summary statement.*** Dr. Castranova concluded that the proposed protocol is a significant advance in asbestos risk assessment methodology. He strongly supported the recommendation that future measurements be performed using TEM, rather than PCM. Dr. Castranova also supported the approach of assigning equal carcinogenic potency to cleavage fragments and asbestos fibers of similar dimension—a finding, he noted, that could be tested in an animal inhalation study. Further, Dr. Castranova agreed that non-asbestiform amphiboles and asbestos amphiboles of the same dimension should be assigned equal carcinogenic potency. Dr. Castranova indicated that the epidemiology and toxicology literature clearly indicate that

mesothelioma potency varies with fiber type, but he was not convinced that this literature supports a difference in lung cancer potency between amphibole and chrysotile fibers.

- # ***Dr. Price's summary statement.*** Dr. Price found the proposed protocol to be an impressive compilation of the epidemiology and toxicology literature into a cancer risk assessment model that addresses most, but not all, risk factors debated since EPA's 1986 model. Dr. Price urged EPA to explore exposure-response models other than the models that involve linear, low-dose extrapolations, which he viewed as being inconsistent with the epidemiology literature. Dr. Price indicated that future revisions to the protocol should definitely consider non-linear models and threshold effects.

As an additional comment, Dr. Price emphasized that the two main elements of the protocol—the proposed exposure index and the exposure-response analysis—are closely inter-related and subsequent changes to the proposed exposure index could affect the robustness of the overall modeling effort. As an example of his concern, Dr. Price noted that increasing the fiber diameter cut-off in the exposure index from 0.5 μm to 1.5 μm could (according to an observer comment) lead to dramatic differences in the number of cleavage fragments counted in environment samples; however, he indicated that the animal studies used to derive the original exposure index did not include cleavage fragments. Such scenarios raise questions about using an exposure index derived from very specific exposure conditions in animal studies to evaluate human health risks associated with exposures of an entirely different character. Dr. Price encouraged further study of cleavage fragments, perhaps in an animal inhalation study, to resolve the role of cleavage fragments.

- # ***Dr. Case's summary statement.*** Dr. Case congratulated Drs. Berman and Crump for compiling what he viewed as a reasonable evaluation of the available toxicology and epidemiology literature, and he strongly supported the general approach of factoring fiber type and fiber dimension into cancer risk assessment. Dr. Case indicated that he agreed with the finding that amphibole fibers have slightly greater lung cancer potency than do chrysotile fibers, although he believed that fiber dose, fiber length, and especially smoking history and type of industry have greater importance in this regard. Dr. Case recognized that how one views the differences between the Quebec and South Carolina cohorts affects the conclusions drawn on this issue, and he encouraged EPA to classify the cohort of South Carolina textile workers as being exposed to mixed asbestos fibers, rather than being exposed to only chrysotile fibers.²

² When presenting the summary statements, one panelist (LS) indicated that NIOSH is re-analyzing filters that were collected in the 1960s from the South Carolina textile plant, and these re-analyses should indicate the distribution of fiber types in this cohort's exposures. Another panelist (BC) noted that these re-analyses will not characterize earlier exposures to amosite fibers, which are believed to have occurred primarily before 1950 (based on findings from studies of lung-retained fibers).

Dr. Case made several recommendations for further evaluating the existing epidemiological data and for collecting additional data. First, Dr. Case indicated that it is critically important for any lung cancer risk model to consider confounding effects of cigarette smoking, and he encouraged EPA to incorporate interactions with cigarette smoking into the lung cancer model to the greatest extent possible. Second, Dr. Case supported Dr. Lippmann's recommendation of convening an additional expert panel workshop to critically review inferences that should be drawn from the exposure measurements made in the epidemiological studies; such a panel, Dr. Case noted, would require inputs from experts in mineralogy, industrial hygiene, and measurement methodologies. Third, he supported comments recommending that EPA examine non-linear and threshold exposure-response models. Finally, Dr. Case agreed that conducting an animal inhalation study is probably the best way to examine whether tremolite cleavage fragments produce lung cancer, but did not advocate using rat inhalation studies to examine whether these fragments induce mesothelioma, because results from rat inhalation studies have been shown to be a poor model for mesothelioma in humans. He added, however, that it would quite probably be impossible to design an experiment in which rats were exposed only to "cleavage fragments" or "non-asbestiform fibers" with no asbestiform fibers present at all.

Dr. Stayner's summary statement. Dr. Stayner supported the general concept of incorporating fiber type and fiber dimension into cancer risk assessment, but he recommended that additional work be conducted before EPA accepts the proposed protocol as a new risk assessment paradigm. Dr. Stayner indicated that his confidence in the proposed protocol varies between the lung cancer and mesothelioma models.

For lung cancer, Dr. Stayner indicated that the available epidemiological data should be able to support a new risk assessment model, but he recommended that EPA consider the panelists' many recommendations for how the meta-analysis can be improved (e.g., using different statistical models, developing and applying minimal study inclusion criteria, conducting additional sensitivity analyses). Concurring with Dr. Steenland's summary statement, Dr. Stayner added that cigarette smoking is very unlikely to be a confounding factor in the lung cancer model and he questioned whether the available data would support a quantitative assessment of the interaction effects. While Dr. Stayner supported the recommendation for evaluating non-linear exposure-response models, he noted that the individual-level data needed to construct these models are not available for most epidemiological studies. Dr. Stayner added that obtaining raw data from additional occupational cohorts would provide the best opportunity for more detailed exploration of non-linear exposure-response relationships.

Dr. Stayner expressed greater concern about the foundation of the mesothelioma risk model. He indicated, for instance, that the relative potencies included in the proposed exposure index are based entirely on toxicology studies for lung cancer, and not on any epidemiology or toxicology studies specific to mesothelioma. Despite these concerns about the biological basis for the proposed mesothelioma model, Dr. Stayner noted that the proposed model does provide an

improved fit to the findings from the epidemiological studies. He recommended that EPA consider optimizing the relative potencies in the exposure index to the human data, especially if EPA can access raw data from additional occupational cohorts to evaluate how exposure-response varies with fiber size and fiber type.

Dr. McClellan's summary statement. Dr. McClellan congratulated Drs. Berman and Crump for integrating the toxicological and epidemiological data into a reasonable evaluation of asbestos cancer risks. Overall, Dr. McClellan found the proposed protocol to be a substantial improvement over EPA's 1986 models and urged EPA to continue to move forward with completing the protocol based on the panelists' feedback. Though he found the presentation of information in the draft document to lack transparency on many important matters, Dr. McClellan indicated that the authors' presentations at the workshop addressed many of his concerns regarding the transparency of how the proposed model was developed. One suggested improvement to the protocol's transparency was to clearly describe what literature were reviewed and to specify what studies actually factored into the quantitative analyses.

Addressing specific topics, Dr. McClellan indicated that the analyses in the proposed protocol adequately characterize the general roles that fiber type and fiber dimension play in cancer risk. He supported suggestions for involving additional experts, perhaps in another expert panel review, to further review interpretations of the epidemiological studies. Further, Dr. McClellan agreed with other panelists' recommendation that EPA explore the utility of non-linear exposure-response models, consistent with the agency's proposed revised Cancer Risk Assessment Guidelines. If linear, low-dose extrapolation models are ultimately used, he suggested that EPA explicitly acknowledge the uncertainties associated with such an approach. Dr. McClellan indicated that obtaining raw data from additional epidemiological studies might be particularly helpful in the exposure-response modeling. Finally, Dr. McClellan emphasized that the exposure characterization in the proposed protocol is closely linked to the exposure-response assessment; thus, the authors must carefully consider how revisions to the exposure characterization affect the assumptions in the exposure-response assessment, and vice versa.

6.2 Development of Final Conclusions and Recommendations

After presenting their individual conclusions and recommendations, the panelists worked together to draft summary statements for the peer consultation workshop. Every panelist was asked to write a brief synopsis of a particular topic debated during the workshop. These draft statements were then displayed to the entire panel and observers, edited by the panelists, and then compiled into this document's

executive summary, which should be viewed as the expert panel's final conclusions and recommendations regarding the proposed protocol.

7. REFERENCES

B Bellmann, H Muhle, O Creutzenberg, et al. 2001. Effects of nonfibrous particles on ceramic fiber (RCF1) toxicity in rats. *Inhalation Toxicology* 13(10):877–901.

DW Berman, KS Crump, EJ Chatfield, JMG Davis, AD Jones. 1995. The Sizes, Shapes, and Mineralogy of Asbestos Structures that Induce Lung Tumors or Mesothelioma in AF/HAN Rats Following Inhalation. *Risk Analysis* 15(2).

DW Berman and KS Crump 1999. Methodology for Conducting Risk Assessments at Asbestos Superfund Sites; Part 1: Protocol. Final Draft. Prepared for U.S. Environmental Protection Agency. February 15, 1999.

DW Berman and KS Crump 2001. Technical Support Document for a Protocol to Assess Asbestos-Related Risk. Final Draft. Prepared for U.S. Environmental Protection Agency and U.S. Department of Transportation. September 4, 2001.

G Berry and ML Newhouse. 1983. Mortality of Workers Manufacturing Friction Materials Using Asbestos. *British Journal of Industrial Medicine* 40:1–7.

C Boutin, P Dumortier, F Rey, et al. 1996. Black spots concentrate oncogenic asbestos fibers in the parietal pleura. *American Journal of Respiratory Critical Care and Medicine* 153:444–449.

M Camus, J Siemiatycki, B Meek. 1998. Nonoccupational exposure to chrysotile asbestos and the risk of lung cancer. *New England Journal of Medicine* 338:1565–1571.

WC Cooper, O Wong, and R Graebner. 1988. Mortality of workers in two Minnesota taconite mining and milling operations. *Journal of Occupational Medicine* 30(6):506–511.

JMG Davis, J Addison, C McIntosh, BG Miller, and K Niven. 1991. Variations in the Carcinogenicity of Tremolite Dust Samples of Differing Morphology. *Annals of the New York Academy of Sciences*, 473–490.

PE Enterline, J Harley, and V Henderson. 1986. Asbestos and Cancer—A Cohort Followed to Death. Graduate School of Public Health, University of Pittsburgh.

N De Klerk, B Armstrong, A Musk, M Hobbs. 1989. Cancer mortality in relation to measures of exposure to crocidolite at Wittenoom Gorge in Western Australia. *British Journal of Industrial Medicine* 46:529–536.

JM Dement, DP Brown, A Okun. 1994. Follow-up Study of Chrysotile Asbestos Textile Workers: Cohort Mortality and Case-Control Analysis. *American Journal of Industrial Medicine* 26:431–447.

EPA 1986. Airborne Asbestos Health Assessment Update. U.S. Environmental Protection Agency. EPA 600/8-84-003F. 1986.

ERG. 2003. Report on the Expert Panel on Health Effects of Asbestos and Synthetic Vitreous Fibers: The Influence of Fiber Length. Prepared by Eastern Research Group, Inc., for the Agency for Toxic Substances and Disease Registry. March 17, 2003.

TW Hesterberg, GA Hart, J Chevalier, et al. 1998. The importance of fiber biopersistence and lung dose in determining the chronic inhalation effects of X607, RCF1, and chrysotile asbestos in rats. *Toxicology and Applied Pharmacology* 153:68–82.

ITT Higgins, JH Glassman, MS Oh, and RG Cornell. 1983. Mortality of reserve mining company employees in relation to taconite dust exposure. *American Journal of Epidemiology* 118(5):710–719.

J Hodgson and A Darnton. 2000. The Quantitative Risk of Mesothelioma and Lung Cancer in Relation to Asbestos Exposure. *Annals of Occupational Hygiene* 44(8):565–201.

JM Hughes, H Weill, YY Hammad. 1987. Mortality of Workers Employed at Two Asbestos Cement Plants. *British Journal of Industrial Medicine* 44:161–174.

IARC. 1996. AB Kane, P Boffetta, R Saracci, and JD Wilbourn. Mechanisms of Fibre Carcinogenesis. International Agency for Research on Cancer. Oxford University Press 140:1–9.

JL Levin, JW McLarty, GA Hurst, AN Smith, and AL Frank. 1998. Tyler Asbestos Workers: Mortality Experience in a Cohort Exposed to Amosite. *Occupational and Environmental Medicine* 55:155–160.

FD Liddell and BG Armstrong. 2002. The combination of effects on lung cancer of cigarette smoking and exposure in Quebec chrysotile miners and millers. *Annals of Occupational Hygiene* 46(1):5–13.

FD Liddell, AD McDonald, and JC McDonald. 1997. The 1891-1920 Birth Cohort of Quebec Chrysotile Miners and Millers: Development from 1904 and Mortality to 1992. *Annals of Occupational Hygiene* 41:13–36.

FD Liddell, AD McDonald, and JC McDonald. 1998. Dust exposure and lung cancer in Quebec chrysotile miners and millers. *Annals of Occupational Hygiene* 42(1):7–20.

M Lippmann. 1988. Review: Asbestos exposure indices. *Environmental Res* 46:86–106.

- M Lippmann. 1994. Deposition and retention of fibers: Effects on incidence of lung cancer and mesothelioma. *Occupational and Environmental Medicine* 51:793–798.
- JC McDonald, GW Gibbs, FD Liddell, and AD McDonald. 1978. Morality after long exposure to cummingtonite-grunerite. *American Review of Respiratory Disease* 118(2):271–277.
- AD McDonald, JS Fry, AJ Woolley, and JC McDonald. 1984. Dust Exposure and Mortality in an American Chrysotile Asbestos Friction Products Plant. *British Journal of Industrial Medicine* 41:151–157.
- JC McDonald, B Armstrong, B Case, et al. 1989. Mesothelioma and asbestos fiber type. Evidence from lung tissue analysis. *Cancer* 63:1544–1547.
- AD McDonald, BW Case, A Churg, A Dufresne, GW Gibbs, P Sebastien, and JC McDonald. 1997. Mesothelioma in Quebec chrysotile miners and millers: epidemiology and aetiology. *Annals of Occupational Hygiene* 41(6):707–719.
- JC McDonald, J Harris, and B Armstrong. 2002. Cohort mortality study of vermiculite miners exposed to fibrous tremolite: an update. *Annals of Occupational Hygiene* 46(S1):93–94.
- WJ Nicholson. 1976. Part III: Recent Approaches to the Control of Carcinogenic Exposures. Case Study 1: Asbestos—The TLV Approach. *Annals of New York Academy of Science* 271:152–169.
- C-G Ohlson, T Rydman, L Sundell, et al. 1984. Decreased lung function in long-term asbestos cement workers: A cross-sectional study. *American Journal of Industrial Medicine* 5:359–366.
- K Rödelsperger, H-J Weitowitz, B Brückel, et al. 1999. Dose-response relationship between amphibole fiber lung burden and mesothelioma. *Cancer Detection and Prevention* 23(3):183–193.
- AJ Rogers, J Leigh, G Berry, et al. 1991. Relationship between lung asbestos fiber type and concentration and relative risk of mesothelioma. *Cancer* 67:1912–1920.
- A Rogers and G Major. 2002. Letter to the Editor: The Quantitative Risks of Mesothelioma and Lung Cancer in Relation to Asbestos Exposure: The Wittenoom Data. *Annals of Occupational Hygiene* 46(1):127–129.
- H Seidman, IJ Selikoff, and SK Gelb. 1986. Mortality Experience of Amosite Asbestos Factory Workers: Dose-Response Relationships 5 to 40 Years after onset of Short-Term Work Exposure. *American Journal of Industrial Medicine* 10:479–514.

IJ Selikoff and H Seidman. 1991. Asbestos-Associated Deaths among Insulation Workers in the United States and Canada, 1967–1987. *Annals of the New York Academy of Sciences* 643:1–14.

LT Stayner, DA Dankovic, RA Lemen. 1996. Occupational Exposures to Chrysotile Asbestos and Cancer Risk: A Review of the Amphibole Hypothesis. *American Journal of Public Health* 86(2):176–186.

V Timbrell, T Ashcroft, B Goldstein, et al. 1988. Relationships between retained amphibole fibers and fibrosis in human lung tissue specimens. In: *Inhaled Particles VI. Annals of Occupational Hygiene* 32(S1):323–340.

Y Suzuki and S Yuen. 2001. Asbestos Tissue Burden Study on Human Malignant Mesothelioma. *Industrial Health* 39:150–160.

MJ Teta, HC Lewinsohn, JW Meigs, et al. 1983. Mesothelioma in Connecticut: 1955–1977: Occupational and geographic associations. *Journal of Occupational Medicine* 25(10):749–756.

A Tossavainen, M Kotilainen, K Takahashi, G Pan, and E Vanhala. 2001. Amphibole Fibers in Chinese Chrysotile Asbestos. *Annals of Occupational Hygiene* 45(2):145-152.

H Weill, JL Abraham, JR Balmes, B Case, AM Churg, J Hughes, M Schenker, and P Sebastien. 1990. Health Effects of Tremolite. Official statement of the American Thoracic Society. *American Review of Respiratory Disease* 142(6):1453–1458.

E Yano, ZM Wang, XR Wang, MZ Wang, and YJ Lan. 2001. Cancer Mortality among Workers Exposed to Amphibole-free Chrysotile Asbestos. *American Journal of Epidemiology* 154(6):538–543.

Appendix A

List of Expert Panelists

Appendix B

Premeeting Comments, Alphabetized by Author (includes bios of panelists and the charge to the panelists)

Note: This appendix is a copy of the booklet of the premeeting comments that ERG distributed at the peer consultation workshop. One panelist (Dr. Bruce Case) submitted an edited form of his premeeting comments to ERG at the workshop. That edited version appears in this appendix.

Appendix C

List of Registered Observers of the Peer Consultation Workshop

Appendix D

Agenda for the Peer Consultation Workshop

Appendix E

Observer Comments Provided at the Peer Consultation Workshop

Note: The peer consultation workshop included three observer comment periods, one on the first day of the workshop and two on the second day of the workshop. This appendix includes verbatim transcripts (to the extent that specific remarks were audible from recordings) of the observer comments, in the order the comments were given.

Appendix F

Observer Post-Meeting Comments